

CORRECTION

Open Access



Correction: Paradoxical tuberculosis-associated immune reconstitution inflammatory syndrome in initiating ART among HIV-Infected patients in China-risk factors and management

Honghong Yang¹, Qian Liu¹, Yushan Wu¹, Kun He¹, Qin Zeng¹ and Min Liu^{1*}

Correction: *BMC Infect Dis* 24, 5 (2024)
<https://doi.org/10.1186/s12879-023-08897-3>

Following publication of the original article [1], we have been notified that Fig. 2 was showing one unnecessary parameter (fever).

The original article can be found online at <https://doi.org/10.1186/s12879-023-08897-3>.

*Correspondence:


Min Liu
gwzxlumin@foxmail.com

¹ Division of Infectious Diseases, Chongqing Public Health Medical Center, 109 Baoyu Road, Shapingba District, Chongqing 400036, China



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Originally published Fig. 2:



Corrected Fig. 2:




Fig. 2 Associations between pre-ART clinical and laboratory characteristics with subsequent paradoxical TB-IRIS events. The variables that showed a significant relationship with the development of paradoxical TB-IRIS (from Table 1), that is, age, haemoglobin, baseline CD4 + T-cell counts, HIV VL, and miliary were included in this predictive model. Association of all variables with risk for TB-IRIS was assessed in adjusted logistic regression models. Odds ratios for values below or above threshold levels were displayed in a forest plot R—odds ratio; CI—confidence level

The original article has been corrected.

Published online: 01 March 2024

Reference

1. Yang H, et al. Paradoxical tuberculosis-associated immune reconstitution inflammatory syndrome in initiating ART among HIV-Infected patients in China-risk factors and management. *BMC Infect Dis.* 2024;24:5. <https://doi.org/10.1186/s12879-023-08897-3>.