

Research article

Open Access

Nosocomial outbreak of neonatal *Salmonella enterica* serotype Enteritidis meningitis in a rural hospital in northern Tanzania

Hogne Vaagland*^{1,2}, Bjørn Blomberg^{2,3}, Carsten Krüger^{4,5}, Naftali Naman⁴, Roland Jureen³ and Nina Langeland^{3,6}

Address: ¹Department of Otolaryngology/Head & Neck Surgery, Haukeland University Hospital, N-5021 Bergen, Norway, ²Centre for International Health, University of Bergen, N-5021 Bergen, Norway, ³Institute of Medicine, University of Bergen, N-5021 Bergen, Norway, ⁴Haydom Lutheran Hospital, Mbulu, Tanzania, ⁵Deipe Stegge 71, D-48653 Coesfeld, Germany and ⁶Department of Medicine, Haukeland University Hospital, N-5021 Bergen, Norway

Email: Hogne Vaagland* - hogne.vaagland@cih.uib.no; Bjørn Blomberg - bjorn.blomberg@cih.uib.no; Carsten Krüger - thea.carsten.krueger@web.de; Naftali Naman - nbnaman@elct.org; Roland Jureen - roland.jureen@med.uib.no; Nina Langeland - nina.langeland@helse-bergen.no

* Corresponding author

Published: 14 September 2004

Received: 27 June 2004

BMC Infectious Diseases 2004, 4:35 doi:10.1186/1471-2334-4-35

Accepted: 14 September 2004

This article is available from: <http://www.biomedcentral.com/1471-2334/4/35>

© 2004 Vaagland et al; licensee BioMed Central Ltd.

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Clinicians at Haydom Lutheran Hospital, a rural hospital in northern Tanzania noted an unusually high case-fatality rate of pediatric meningitis and suspected an outbreak of an unknown agent or an organism resistant to the empirical therapy.

Methods: We established a provisional microbiology laboratory to investigate the suspected outbreak. Blood and spinal fluid specimens were taken from children below the age of seven years with suspected meningitis. The blood and spinal fluid specimens were inoculated in commercial blood culture bottles and locally prepared Thayer-Martin medium in slanted tubes, respectively. The bacterial isolates were sent to Norway for further investigation, including susceptibility testing and pulsed-field gel-electrophoresis (PFGE).

Results: Among 24 children with suspected meningitis and/or septicemia, five neonates had meningitis caused by *Salmonella enterica* serotype Enteritidis, all of whom died. Two children had *S. Enteritidis* septicemia without meningitis and both survived. Genotyping with PFGE suggested a clonal outbreak. The salmonella strain was resistant to ampicillin and sensitive to gentamicin, the two drugs commonly used to treat neonatal meningitis at the hospital.

Conclusion: The investigation reminds us that nontyphoidal salmonellae can cause meningitis associated with very high case-fatality rates. Resistance to multiple antimicrobial agents increases the risk of treatment failure and may have contributed to the fatal outcome in all of the five patients with salmonella meningitis. The investigation indicated that the outbreak was nosocomial and the outbreak subsided after hygienic measures were instituted. Establishing a provisional microbiological laboratory is a valuable and affordable tool to investigate and control outbreaks even in remote rural areas.

Background

Nontyphoidal salmonellae are a common cause of food-borne illnesses. In Africa, nontyphoidal salmonellae are the most common cause of bloodstream infections in children younger than five years [1]. While meningitis caused by nontyphoidal salmonellae is uncommon in economically developed countries [2], it is more frequent in tropical countries, particularly in children younger than six months, and associated with higher case-fatality rates than meningitis caused by other bacteria [1,3-5].

At Haydom Lutheran Hospital, a rural hospital in northern Tanzania, clinicians noted an extraordinarily high case-fatality rate (>60%) from pediatric meningitis in the period January 1998 to April 2000. It was suspected that the high case-fatality rate of meningitis was due to an outbreak of an unusual etiological agent, or an organism resistant to the hospital's standard empirical treatment, which was ampicillin and gentamicin for infants (<2 months) and penicillin and chloramphenicol for older children. Thus, we established a provisional microbiology laboratory to identify the causative agents and to facilitate the implementation of effective preventive measures.

Methods

From a total of 360 children admitted from July to August 2000, blood and/or spinal fluid specimens were collected from 24 children aged one day to six years (median age 23.5 days) with suspected meningitis and/or septicaemia, after careful evaluation by the attending pediatrician. Blood and spinal fluid specimens were inoculated in BBL SeptiChek blood-culture bottles (Becton Dickinson, Sparks, MD USA) and on locally prepared non-selective Thayer-Martin medium in slanted tubes, respectively. All cultures were incubated at 35°C for 5 days and inspected daily for bacterial growth. Positive bacterial specimens were shipped to Institute of Medicine, Haukeland University Hospital, Norway, for further study. The total cost of the laboratory reagents used on site was \$475. Positive specimens were identified using standard laboratory methods [6]. The susceptibilities of the isolates to antimicrobial agents were examined by disk diffusion method on PDM medium (AB Biodisk, Solna, Sweden) [7]. The isolates were genotyped with pulsed-field gel electrophoresis (PFGE). Statistics were calculated with Stata 8 for Mac OSX (Stata Corporation, College Station, TX).

Results

On clinical grounds, twenty-four children were considered to have possible sepsis or meningitis. Blood culture was taken from 23 children, but 13 had received prior antibiotic treatment. Spinal fluid culture was taken from 16 children, of whom ten had received prior antibiotic treatment. Both blood culture and spinal fluid culture was obtained from a total of 15 children. *Salmonella enterica*

serotype Enteritidis was isolated from seven patients, of whom four had positive cultures from both blood and spinal fluid and three from blood only. One of the three patients, who had *S. Enteritidis* isolated only from blood culture, had pus in the spinal fluid, and was therefore considered a case of *S. Enteritidis* meningitis, resulting in a total of five cases of *S. Enteritidis* meningitis. One isolate each of *Staphylococcus aureus* and *Streptococcus pyogenes* were isolated from spinal fluid and three isolates of coagulase-negative staphylococci were isolated from blood. Antimicrobial susceptibility testing showed that all the *S. Enteritidis* isolates were resistant to chloramphenicol and ampicillin, intermediate resistant to cefuroxime, and sensitive to gentamicin, cefotaxime and ciprofloxacin. As shown in Figure 1, the PFGE patterns were identical for all the eleven *S. Enteritidis* isolates except one, which differed by only one band (Figure 1 shows a total of 17 strains, 6 of which are duplicate isolates from the same patients). All children with *S. Enteritidis* infections were neonates (median age 15 days, range: 8 to 27 days), whereas those with other infections or negative cultures, on average, were older (median age 4 months, range: 1 day to 6 years). All children with *S. Enteritidis* infections had been delivered at Haydom Lutheran Hospital, and five out of the seven never left the hospital before they became ill. Table 1 shows the characteristics of the patients with *S. Enteritidis* infection. All five children with *S. Enteritidis* meningitis died. The two children with *S. Enteritidis* sepsis survived. Five (29%) of the 17 children without verified *S. Enteritidis* infection also died. Among the 24 children investigated, *S. Enteritidis* meningitis was associated with a relative risk of 3.8 (95% confidence interval 1.8 to 8.1) for fatal outcome.

Discussion

The finding of an outbreak of bacteremia and meningitis caused by *S. Enteritidis* was not anticipated. Genotyping with PFGE suggested a clonal outbreak. This genotyping information, the susceptibility patterns and the clinical information that all children with *S. Enteritidis* infections were born at the hospital and that the majority never left the hospital before they became ill, strongly suggest that the outbreak was nosocomial. Nontyphoidal salmonella infections are frequently associated with animal reservoirs and infection usually originates from food products. Nosocomial spread of nontyphoidal salmonellae, particularly in neonatal wards, is known from the literature [8]. Neonates are at particular risk of infection because of relatively reduced gastric acidity and peristalsis [1]. In previous nosocomial outbreaks caused by salmonellae, the sources of infection have been related to the use of contaminated medications, diagnostics, blood products, banked human milk, the use of raw eggs or yeast in tube feeding and improper disinfection of devices such as rubber tubes for oropharyngeal suction [8,9]. The exact




Figure 1

Pulsed-field gel-electrophoresis (PFGE) of digested DNA from 17 isolates of *Salmonella enterica* serotype Enteritidis obtained from seven children with meningitis and/or septicemia. The isolates were obtained from blood cultures of all seven children and spinal fluid culture of four children. Six of the isolates in the gel are duplicate isolates from the same children.

source of the outbreak at the hospital was not established. Direct food-borne transmission was unlikely, since the neonates at the hospital were fed on breast milk. Possible sources include contaminated instruments, clothes or bathing facilities for the newborn. Spread of infection may have occurred by baby-to-baby transmission or via family members and/or hospital staff. The medical staff at the hospital was informed about the findings, and immediate interventions in the form of hygienic measures were instituted, including the reinforcement of disinfection and hand-washing practices. Data from the hospital annual reports shows that the case-fatality rate from pediatric meningitis dropped from >60% before the intervention to 40% by 2001.

The *S. Enteritidis* strain responsible for the outbreak was resistant to two of the first-line drugs, ampicillin and chloramphenicol, but sensitive to gentamicin. This finding is not surprising, since multi-drug-resistant *S. Enteritidis* has been reported from this region for decades [10]. The children at the hospital were treated with a combination of ampicillin and gentamicin. The high case-fatality rate in these patients implies that *de facto* monotherapy with gentamicin may be suboptimal as treatment for *S. Enteritidis* meningitis considering that gentamicin traverses the blood-brain-barrier poorly, is bound to proteins and inhibited by the acidity in infected cerebrospinal fluid. However, it also reflects the poor prognosis of neonatal *S. Enteritidis* meningitis regardless of therapy. Molyneux

Table 1: Characteristics of neonates with meningitis (n = 5) and sepsis (n = 2) caused by *Salmonella enterica* serovar Enteritidis at Haydom Lutheran Hospital, Northern Tanzania, in July and August 2000

Patient characteristic	Pat. 1	Pat. 2	Pat. 3	Pat. 4	Pat. 5	Pat. 6	Pat. 7
Sex	F	F	M	F	F	F	M
Mode of delivery	SVD	SVD	Sectio	SVD	SVD	SVD	SVD
Age at onset	8 d	7 d	8 d	15 d	24 d	14 d	13 d
Age at discharge (*death)	20 d	21 d	32 d*	21 d*	29 d*	28 d*	17 d*
Birth weight	2350 g	4000 g	3300 g	2620 g	3600 g	ND	2800 g
Fever	+	-	+	+	+	+	-
Neck stiffness	-	-	-	-	-	-	-
Fits	-	-	-	+	+	+	+
Bulging fontanel	-	-	-	+	+	-	-
<i>S. Enteritidis</i> in blood culture	+	+	+	+	+	+	+
<i>S. Enteritidis</i> in CSF culture	ND	-	ND	+	+	+	+
CSF WBC	ND	10/ μ L	Pus	Pus	>3000/ μ L	ND	32/ μ L
Clinical diagnosis							
- Meningitis	-	-	+	+	+	+	+
- Septicemia	+	+	+	-	-	+	+
- Malaria	+	+	-	-	+	-	-
Outcome	Alive	Alive	Death	Death	Death	Death	Death

Pat. = Patient, F = Female, M = Male, SVD = Spontaneous vaginal delivery, d = Days, "+" = Present, "-" = Absent, CSF = Cerebrospinal fluid, ND = No data, WBC = White blood cells

reported fatal outcome for 58% of children with meningitis caused by nontyphoidal salmonellae despite routine treatment with chloramphenicol, to which all bacterial isolates were sensitive in vitro [4]. Many authorities recommend third-generation cephalosporins as empirical chemotherapy for meningitis caused by gram-negative bacteria [11], not only because of high bactericidal activity due to low minimum inhibitory concentrations (MICs), but also because they penetrate the blood-brain-barrier better than both gentamicin and chloramphenicol. A third-generation cephalosporin, such as cefotaxime would have been an excellent therapeutic option in this case. However, the price of the newer cephalosporins is often prohibitive in the setting of low-income countries such as Tanzania. Ciprofloxacin is generally not recommended for use in children due to potential adverse effects, but can be resorted to for treatment of life-threatening infections with multidrug-resistant nontyphoidal salmonellae [12]. The standard empirical treatment regimen at the hospital could not be changed due to financial constraints.

Infections caused by nontyphoidal salmonellae in children in Africa are more common during the rainy season and have been associated with malaria, anemia and malnutrition [1]. However, there is considerable overlap between these medical conditions, all of which may be more frequent during the rainy season. This investigation was performed during the dry season. Infections caused by nontyphoidal salmonellae have also been associated with HIV infection [1]. We do not know the HIV status of

the children involved in this outbreak, however, in this part of Tanzania, the rate of HIV infection is less than 0.5% [13].

Conclusions

There are a number of lessons to be learned from this investigation. We are reminded that *S. Enteritidis* can cause meningitis, which carries a very high case-fatality rate [1,3,4]. Our findings support former reports that *S. Enteritidis* can easily spread by nosocomial transmission, particularly in neonatal wards. Resistance to multiple antimicrobial agents increases the risk that empirical therapy will fail, especially in settings where modern cephalosporins are not affordable. Without adequate laboratory facilities, correct diagnosis and treatment of bacterial meningitis and septicemia in children remains a challenge. However, the report also shows that establishing a provisional microbiology laboratory can be a valuable and affordable tool to investigate and curb epidemics even in the setting of remote rural Africa, provided there is a proficient laboratory willing to assist.

Competing interests

None declared.

Authors' contributions

HV was the principal investigator, participated in the planning and execution of the work, including performing the on-site microbiological procedures, and analysis of data, and was the main responsible author. BB participated in planning, microbiological investigations in Nor-

way, data analysis and writing. CK and NN undertook clinical investigation and sample collection and participated in the writing. RJ was responsible for the microbiological investigations in Norway and participated in the data analysis and writing. NL was the project coordinator and participated in planning, data analysis and writing. All authors read and approved the final manuscript.

Acknowledgements

We thank all nursing and medical staff at Haydom Lutheran Hospital who cared for the children and facilitated this work and Dr Bjørg Evjen Olsen, who provided important information, including the relevant hospital annual reports. We are particularly grateful to the director of the hospital Dr Ole Hallgrim Evjen Olsen for his willingness and interest in having this investigation undertaken.

References

- Graham SM, Molyneux EM, Walsh AL, Cheesbrough JS, Molyneux ME, Hart CA: **Nontyphoidal Salmonella infections of children in tropical Africa.** *Pediatr Infect Dis J* 2000, **19**:1189-1196.
- Saphra I, Winter JW: **Clinical manifestations of salmonellosis in man; an evaluation of 7779 human infections identified at the New York Salmonella Center.** *N Engl J Med* 1957, **256**:1128-1134.
- Barclay N: **High frequency of salmonella species as a cause of neonatal meningitis in Ibadan, Nigeria. A review of thirty-eight cases.** *Acta Paediatr Scand* 1971, **60**:540-544.
- Molyneux EM, Walsh AL, Malenga G, Rogerson S, Molyneux ME: **Salmonella meningitis in children in Blantyre, Malawi, 1996-1999.** *Ann Trop Paediatr* 2000, **20**:41-44.
- Sirinavin S, Chiemchanya S, Vorachit M: **Systemic nontyphoidal Salmonella infection in normal infants in Thailand.** *Pediatr Infect Dis J* 2001, **20**:581-587.
- Murray PR, Baron EJ, Jorgensen JH, Pfaller MA, Tenover FC, Tenover FC: **Manual of Clinical Microbiology.** 8th edition. Washington D. C., ASM Press; 2003.
- Bergan T, Bruun JN, Digraanes A, Lingaas E, Melby KK, Sander J: **Susceptibility testing of bacteria and fungi. Report from "the Norwegian Working Group on Antibiotics".** *Scand J Infect Dis Suppl* 1997, **103**:1-36.
- Khan MA, Abdur-Rab M, Israr N, Ilyas M, Ahmad F, Kundi Z, Ghafoor A: **Transmission of Salmonella worthington by oropharyngeal suction in hospital neonatal unit.** *Pediatr Infect Dis J* 1991, **10**:668-672.
- Weikel CS, Guerrant RL: **Nosocomial salmonellosis.** *Infect Control* 1985, **6**:218-220.
- Petat E, Carteron B, Reguer M, Lemmens P, Vandepitte J, Ghysels G: **Étude des Shigella et Salmonella isolées au Burundi de 1980 à 1985.** *Bull Soc Pathol Exot Filiales* 1987, **80**:171-179.
- Quagliarello VJ, Scheld WM: **Treatment of bacterial meningitis.** *N Engl J Med* 1997, **336**:708-716.
- Wessalowski R, Thomas L, Kivit J, Voit T: **Multiple brain abscesses caused by Salmonella enteritidis in a neonate: successful treatment with ciprofloxacin.** *Pediatr Infect Dis J* 1993, **12**:683-688.
- Hinderaker SG, Krüger C, Olsen BE, Naman N, Bergsjø P, Olsen OHE: **Low HIV-seroprevalence in pregnant women in a rural area in Tanzania.** *Acta Obstet Gynecol Scand* 2001, **80**:1152-1153.

Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1471-2334/4/35/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

