

RESEARCH ARTICLE

Open Access



Characteristics and prognostic factors of COVID-19 among infected cases: a nationwide Tunisian analysis

Chahida Harizi^{1,2*}, Ines Cherif^{2,3}, Nourhene Najar^{2,3}, Molka Osman^{2,3}, Rym Mallekh^{2,3}, Oumaima Ben Ayed^{2,3}, Yosr Ayedi^{2,3}, Sonia Dhaouadi^{2,3}, Aicha Hchaichi^{2,3}, Mouna Safer^{2,3}, Hejer Letaief^{2,3}, Ilhem Bouaziz³, Sondes Derouiche^{2,4}, Donia Gharbi^{2,4}, Leila Bouabid^{2,4}, Souha Bougatef^{2,4}, Hamida Ben Salah⁴, Radhouane Fakhfakh^{1,2}, Salma Abid^{2,4}, Ilhem Boutiba Ben Boubaker^{2,4}, Mohamed Kouni Chahed^{2,5} and Nissaf Bouafif Ben-Alaya^{2,3}

Abstract

Background: The outbreak of coronavirus disease (COVID-19) continues to constitute an international public health concern. Few data are available on the duration and prognostic factors of the disease. We aimed to study the recovery time among a Tunisian cohort of COVID-19 confirmed patients and identify the prognostic factors.

Methods: A retrospective, nationwide study was conducted from March 2 to May 8, 2020, recruiting all patients who were diagnosed with COVID-19, by RT-PCR methods, in Tunisia. Data were collected via phone call interview. Kaplan-Meier Methods and Cox proportional hazards regression models were, respectively, used to study the recovery time and estimate its prognostic factors.

Results: One thousand and thirty patients with COVID-19 (aged 43.2 ± 18.2 years, 526 female (51.1%)) were enrolled. Among them 141 (14.8%) were healthcare professionals. Out of 173 patients (17.8%) admitted to the hospital, 47 were admitted in an intensive care unit. Among 827 patients who didn't require specialized care, 55.5% were self-isolated at home, while the rest were in specialized centers. Six hundred and two patients were symptomatic. A total of 634 (61.6%) patients have recovered and 45 (4.4%) patients died. The median duration of illness was estimated to be 31 days (95% CI: [29–32]). Older age (HR = 0.66, CI: [0.46–0.96], $P = 0.031$) and symptoms (HR = 0.61, CI: [0.43–0.81], $P = 0.021$) were independently associated with a delay in recovery time. Being a healthcare professional (HR = 1.52, CI: [1.10–2.08], $P = 0.011$) and patients in home isolation compared to isolation centers (HR = 2.99, CI: [1.85–4.83], $P < 10^{-3}$) were independently associated with faster recovery time.

Conclusion: The duration of illness was estimated to be 1 month. However, this long estimated duration of illness may not equate to infectiousness. A particular attention must to be paid to elderly and symptomatic patients with closer monitoring.

Keywords: COVID-19, SARS CoV-2, Prognostic factors, Clinical characteristics

* Correspondence: chahidaharizi@hotmail.fr

¹Department of Epidemiology and Statistics, Abderrahman Mami Hospital, Ariana, Tunisia

²Faculty of Medicine of Tunis, University of Tunis El Manar, Tunis, Tunisia

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Background

Coronavirus disease 2019 (COVID-19), an infection caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2), was first reported in December 2019, in Wuhan, Hubei Province in China [1]. Since then, human-to-human transmission rates across the world has soared dramatically making it one of the most challenging global health problems. On 30 January 2020, the World Health Organization considered COVID-19 as a public health emergency of international concern and announced the pandemic threat in March 11, 2020 [1, 2]. On May 15, 2020, 4,553,394 confirmed cases of COVID-19 were identified, worldwide, with a death rate of 6.7 and 37.7% of recovery [3].

In Tunisia, the first confirmed case of COVID-19 was reported on March 2, 2020. Since then, several measures have been taken to break the spread of the epidemic among the population. These steps include the successive screening at point of entry, 14 days isolation of travelers returning from risk areas, closure of schools and university facilities, ban of sports and cultural events, borders closure, curfew and finally, a national lockdown was announced on March 22, 2020 [4]. On May 8, 2020, the total number of confirmed cases of COVID-19 was 1030 [5].

Several studies have addressed clinical, radiological and biological issues in COVID-19 patients [6–10]. The clinical spectrum is large and ranges from asymptomatic or mild forms, accounting for about 80%, to cases of acute respiratory distress syndrome that may lead to intensive care or death [11].

However, we note a scarcity of data on prognostic factors associated with a better outcome and recovery. Since an early recovery reduces the shedding and spread of the virus, identifying recovery prognostic factors may improve our understanding of the dynamics of transmission and foster the control of the epidemic [12].

The present study aims to estimate recovery time and identify related prognostic factors among Tunisian infected people with COVID-19.

Methods

Study design and population

This was a retrospective nationwide cohort study. It included all confirmed COVID-19 cases listed among the database of the national observatory of new and emerging diseases (ONMNE), which is the national center for disease control and surveillance, from March 2 to May 8, 2020.

Case definition

A confirmed COVID-19 case was defined as any person, whether symptomatic or not, with laboratory confirmation of SARS-CoV-2 infection using real-time reverse transcriptase–polymerase chain reaction (RT-PCR).

Recovery monitoring and criteria

From March 2, 2020, when the first case was reported, all confirmed cases have been consistently and timely reported by the laboratory to the national observatory for new and emerging diseases (ONMNE).

All positive cases were isolated for at least 14 days, at home or in dedicated COVID-19 centers, and were not released until they got two negative laboratory test results. Hospitalization was only indicated for the serious cases requiring a respiratory assistance or special care. A questionnaire was administered by a team of trained physicians, who were responsible to ensure the follow-up of all confirmed cases by phone calls. This questionnaire included questions about sociodemographic characteristics, isolation setting, symptoms and treatment. The recovery statement was based on laboratory data. For asymptomatic patients, the first recovery testing was scheduled for the last day of the isolation period. For symptomatic patients, the first recovery testing was done 3 days after the symptoms disappearance and at least 7 days after onset of symptoms. In both situations, the second recovery testing was done 24 to 48 h after the first negative sample. If the first recovery test was still positive, then the second recovery test was done 7 days later.

The formal criteria for recovery was defined by two successive negative RT-PCR control specimens and recovery time was defined as the period from the date of symptoms onset to the date of a second negative test. (When patients got a positive RT-PCR result following a false negative result, it was considered as non-effective negative result in our study). After recovery statement, home auto-isolation was advised.

Laboratory confirmation of SARS-CoV-2 infection

The laboratory testing for SARS by RT-PCR was the laboratory method of reference to confirm cases. Charles Nicolle hospital laboratory, which is the national influenza World Health Organization (WHO) collaborating laboratory, was the reference laboratory at national level. It received all the specimens during the first period of outbreak investigation. Later on, other laboratories performing SARS-specific PCR tests were involved.

Detection of SARS-CoV-2 in nasopharyngeal (NP) specimens was performed at the National Reference Laboratory. RNA was extracted from NP swabs using the GXT NA extraction kit (Hainlifescience GmbH, Nehren, Germany) or the QIAamp Viral RNA Mini kit (Qiagen®, Courtaboeuf, France) following the manufacturers' instructions. For SARS-CoV-2 RNA detection, the in-house RT-PCR was used according to the Hong Kong protocol (based on two multiplex assays: the N gene RT-PCR as a screening assay and the Orf1b assay as a confirmatory one). With this assay, a positive COVID-19 result

is determined when both targets (N and Orflab) reach a defined cycle threshold (Ct) of less than 40 [13].

Statistical analysis

Statistical analysis was performed using SPSS version 17.0 software. Data were expressed as frequencies and percentages for categorical variables and as means and standard deviations (SD) or medians and interquartile ranges (IQR) as appropriate for continuous variables. The median time to recovery was estimated by Kaplan-Meier Methods. Cox proportional hazards regression models were used to estimate factors influencing the recovery time, with the hazards ratio (HR) and its 95% confidence interval (95%CI). Statistical significance was determined at $P < 0.05$.

Results

Clinical profile of patients

A total of 1030 patients tested positive for COVID 19 disease were collected between March 2 and May 8, 2020. The mean age was 43.2 years \pm 18.2, and 526 (51.1%) were female. Ninety seven percent of patients were Tunisians (Table 1) and quarter of infections were imported, mainly from France or Turkey. One hundred and seventy three patients (17.8%) were admitted to the hospital, most of them in a non-intensive care department and only 47 in an intensive care unit (ICU). Sixty six patients (38.2%) received an hydroxy-chloroquine treatment (HCQ). Half of the cases received the HCQ in a hospital setting and the others received it either from a private doctor or as self-medication. Patients who did not require care have been self-isolated at home (55.5%) or in dedicated centers for positive cases (44.5%). One hundred and forty-one patients were healthcare professionals. The source of contamination was the healthcare environment in half of the cases.

Seventy five percent of patients had symptoms. The most common symptoms were fatigue, cough and fever, which occurred respectively in 51.6, 51.3 and 48.3% of symptomatic patients. Symptoms such as headache, anosmia and digestive signs were also common in those patients (Table 2).

As of May 8, 2020, a total of 634 (61.6%) patients were recovered, 45 (4.4%) patients were dead and 351 (34.0%) were still being followed up.

Recovery prognostic factors

Survival analysis was used to analyze the prognostic factors of the disease in patients with COVID-19. It showed that the median duration of illness was estimated to be 31 days (95%Confidence Interval [CI]:[29–32])(Fig. 1). In univariate analysis, older age (Fig. 2), imported cases and being symptomatic (Fig. 3) were associated with a longer duration of illness. Whereas, being a healthcare worker

Table 1 Demographic and clinical characteristics of patients with COVID-19 in Tunisia

Characteristics	N (%)
Age groups (years)	
< 15	63 (6.3)
15–34	287 (28.9)
35–54	369 (37.2)
55–74	232 (23.4)
\geq 75	42 (4.2)
Sex	
Male	504 (48.9)
Female	526 (51.1)
Nationality	
Tunisian	999 (97.0)
Non Tunisian	31 (3.0)
Medical staff	
Yes	141 (14.8)
No	809 (85.2)
Contamination mode	
Imported contamination	246 (23.9)
Native cases	783 (76.1)
Hospitalization	
Yes	173 (17.8)
No	800 (82.2)
Admission to ICU	
Yes	47 (4.8)
No	926 (95.2)
Isolation mode	
Home isolation	459 (55.5)
Isolation center	368 (44.5)
Symptoms	
Yes	602 (74.7)
No	204 (25.3)
Chloroquine treatment	
Yes	66 (7.8)
No	779 (92.2)
TOTAL	1030 (100.0)

ICU Intensive care unit

(Fig. 4) and patients in home isolation (Fig. 5) were associated with a shorter duration of illness. Using Cox regression, older age (HR = 0.66, CI:[0.46–0.96], $P = 0.031$) and having symptoms (HR = 0.61, CI:[0.43–0.81], $P = 0.021$) were independently associated with a delay in recovery time. Being a healthcare professional (HR = 1.52, CI: [1.10–2.08], $P = 0.011$) and patients in home isolation compared to isolation centers (HR = 2.99, CI:

Table 2 Distribution of clinical signs among symptomatic patients with COVID-19 in Tunisia

Symptoms	N (%)
Cough	311 (51.6)
Fatigue	309 (51.3)
Fever	291 (48.3)
Headache	210 (34.8)
Anosmia	195 (32.4)
Digestive signs	188 (31.2)
Loss of taste	165 (27.4)
Dyspnea	140 (23.3)
Others signs	129 (21.4)
TOTAL	602 (100.0)

[1.85–4.83], $P < 10^{-3}$) were independently associated with faster recovery time (Table 3).

Discussion

The main result of our study was the long duration of the disease. It took 31 days from the symptoms onset date, for symptomatic patients, or the laboratory confirmation date, for asymptomatic patients, until the second negative sample.

Our national cohort reached 1030 COVID-19 infected patients. It is a relatively young population with most patients under 55 years of age. In the literature, a mean age between 48 and 55 years was reported [6, 14–17]. For the gender, there was no predominance of men or

women. The number of imported cases was higher than the number of native cases until 18th March, 2020. The trend reversal (native cases vs imported cases) was evident from 03/21/2020 which implied an increase in horizontal transmission [5].

Most of cases were mild as evidenced by relatively low ICU admissions and a high number of asymptomatic patients with mild symptoms that do not require treatment in a hospital structure. In fact, the proportion of mild and asymptomatic cases versus severe and fatal cases for COVID-19 infection is currently still unknown. According to a Chinese report publishing all confirmed, suspected, and asymptomatic cases in China, 80% of infections were mild and could recover at home [18]. Concerning clinical presentation of COVID-19 disease, the common symptoms reported by symptomatic patients were cough, fatigue and fever in nearly half of cases and headache in 35% of cases, which was consistent with general symptoms of viral infection. Similar symptoms were described in several case series [6, 15, 19]. One out of three symptomatic patients reported loss of smell and/or taste. This type of disturbance was widely reported in the literature specially among patients with mild and moderate forms of coronavirus disease [20, 21]. A multicenter European study showed that 85.6 and 88.0% of patients reported olfactory and gustatory dysfunctions [22].

The median duration of illness was estimated to be 31 days counting from the day of onset symptoms for symptomatic patients and from confirmation date for






Fig. 1 Survival curve showing survival time to recovery from COVID-19 infection







Table 3 Factors associated with recovery time among patients with COVID-19 in Tunisia

	Univariate Cox regression			Multivariate Cox regression		
	HR	95% CI	P value	HR	95% CI	P value
Male	0.93	0.79–1.09	0.405			
Age > 60 years	0.68	0.55–0.85	0.001*	0.66	0.46–0.96	0.031*
Tunisian Nationality	0.93	0.57–1.50	0.754			
Healthcare worker	1.86	1.52–2.27	< 10 ⁻³ *	1.52	1.10–2.08	0.011*
Imported infection	0.83	0.69–0.99	0.038			
Home isolation	1.69	1.4–2.06	< 10 ⁻³ *	2.99	1.85–4.83	< 10 ⁻³ *
Hospitalization	0.95	0.76–1.18	0.635			
Admission in ICU	0.95	0.71–1.28	0.73			
Symptoms	0.56	0.46–0.68	< 10 ⁻³ *	0.61	0.43–0.81	0.001*
Chloroquine treatment	1.09	0.82–1.46	0.55			

* Significant *p* values

asymptomatic patients. This estimation for illness duration was long. In a recent Singapore study, similar results were recorded. It indicated that by day 15 from onset of illness, only 30% of all COVID-19 patients were PCR-negative by nasopharyngeal swab; this rose to 95% by day 33 [23]. According to these results, the duration of viral shedding may extend to a month and sometimes longer for a small group of patients. Current guideline suggested two consecutive negative RT-PCR test results is one of the criteria for hospital discharge or discontinuation of isolation. However, a high false negative rate of viral test was reported and some patients experienced a “turn positive” of nucleic acid detection by RT-PCR test for SARS-CoV-2 after two consecutive negative results, which may be related to the false negative of RT-PCR test and prolonged nucleic acid conversion [24, 25]. The Singapore study indicates also that viable virus was not found after the second week of illness despite the persistence of PCR detection of RNA [23]. It is, then, important to note that traces of virus detected by RT-PCR were not necessarily correlated with the ability of transmission.

In addition, a recent Canadian study demonstrate that infectivity as defined by growth cell culture were most likely between days one and five [26]. These data indicate that even viral RNA detection may persist in some patients, such persistent RNA detection represent non-viable virus and such patients are non-infectious [23]. These findings are consistent with the new WHO recommendations for discontinue transmission-based precautions (including isolation) and release from the COVID-19 care pathway with 10 days after symptom onset, plus at least 3 days without symptoms for symptomatic patients and 10 days after test positive for asymptomatic patients [27]. These new recommendations should be taken in a count to adapt strategy to discontinue isolation of COVID-19 patients in our country.

Cox regression model showed that younger patients recover faster compared to elderly patients. Similar findings were also reported in literature [28, 29]. In fact, many studies found a high proportion of severe cases and fatality rate among elderly patients with COVID-19 [16, 30–33].

Based on the result of this study, being a healthcare worker was showed to be significantly associated to a lower median time to viral clearance after onset of symptoms. This finding could be explained by the fact that the healthcare workers’ monitoring and control was different from that of other COVID-19 patients and that the deadlines for carrying out the control samples were shorter. Indeed, the COVID-19 monitoring and control protocol adopted by the Tunisian health authorities was inspired from the recovery criteria of the European center for disease prevention and control (ECDC) documents [34]. This protocol consists in carrying out a control sample for asymptomatic patients after 14 days from the date of confirmation. For symptomatic ones, if the fever or any other signs persist beyond 14 days, the control sample was postponed for 3 days after the disappearance of the last sign. These deadlines are not respected for healthcare staff involved in the clinical management of COVID-19 patients. Healthcare workers infected with COVID-19 follow a closer control protocol, which may explain the faster time to viral clearance among them.

Results showed that the place of isolation is playing a significant role in survival time to recovery from COVID-19 infection. According to the output of the cox model, COVID-19 patients in self-isolation at home are almost three times more likely to recover faster than those in dedicated COVID-19 centers. In fact, the Tunisian strategy was based in the first phase of the epidemic on the isolation of confirmed COVID-19 cases at home or in dedicated centers and to hospitalize only serious cases requiring special care. Afterwards, given the registration of an increasing number of secondary cases among

the close contacts of confirmed patients, the health authorities decided to create specialized centers to isolate COVID-19 patients in order to limit community transmission. The delay in recovery for COVID-19 patients isolated in COVID-19 centers compared to those in self-isolation at home could be explained by the fact that patients do not respect self-isolation inside these centers leading sometimes to mass gatherings. This promiscuity inside the dedicated centers for infected people may thus promote the maintenance of viral load. Symptomatic patients seem also to have slower recovery duration.

This study has some limitations. There was a lack in some clinical information such as comorbidities and vital signs (heart rate, respiration rate, blood pressure etc). In addition, since we used phone interview, information concerning, biological and radiological data were not available for patients who have been hospitalized.

Conclusions

Isolation of COVID-19 patients in dedicated centers or at home until the negativation of their second control sample was the strategy that Tunisia followed to combat the novel coronavirus. According to the results of our national study, the estimated duration of illness was extremely long, implying the isolation of confirmed COVID-19 cases for long periods. This partly explains the success in controlling the epidemic in Tunisia. However, with current science data, which proves that viral detection by PCR does not equate to infectiousness, we strongly recommend to lighten our isolation strategy in accordance with the new WHO recommendations.

The results we found reinforce guidance that additional control measures must be undertaken to avoid mass gatherings inside isolation centers. Elderly and symptomatic patients need particular attention and closer monitoring.

Abbreviations

COVID-19: Coronavirus disease; SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2; WHO: World Health Organization; ONMNE: The national observatory of new and emerging diseases; RT-PCR: Real-time reverse transcriptase–polymerase chain reaction; SARS: Severe acute respiratory syndrome; ECDC: The European center for disease prevention and control.

Acknowledgements

We would like to thank the field teams, members of the regional monitoring units and all lab team contributing to the laboratory diagnosis based on the real-time RT-PCR test in particular: Triki Henda, Maha Mastouri, Naila Hannachi and Hakim Hela.

Authors' contributions

CH interpreted the data and drafted the manuscript. IC, NN, MO, RM, OB, YA, [SD]1 and IB collected the data and performed the literature research. AH, MS, HL and RF contributed to data analysis and reviewed multiple drafts of the article. MKC and NBB designed the study, provided guidance and approved the final manuscript. [SD]2, DG, LB, SB and HBS participated in data collection and in the conception of the work. SA and IBB performed laboratory analysis and edited the manuscript. All authors discussed the results, critically read and revised the manuscript and approved its final submitted version.

Funding

The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The follow-up data of individual patients were collected via National Observatory of New and Emerging Diseases of Tunis, Tunisia. Administrative permissions were required to access and use the meta-data described in our study. Permission was granted from the National Observatory of New and Emerging Diseases of Tunis, Tunisia. Data were anonymized for this study. Neither ethical approval nor individual consent was applicable.

Consent for publication

Not applicable.

Competing interests

None declared.

Author details

¹Department of Epidemiology and Statistics, Abderrahman Mami Hospital, Ariana, Tunisia. ²Faculty of Medicine of Tunis, University of Tunis El Manar, Tunis, Tunisia. ³National Observatory of New and Emerging Diseases of Tunisia, Tunis, Tunisia. ⁴Charles Nicolle Hospital, Laboratory of Microbiology, Virology Unit, National Influenza and other Respiratory Viruses Center-Tunisia, Tunis, Tunisia. ⁵Department of Epidemiology and Public Health, Faculty of Medicine of Tunis, Tunis El-Manar University, Tunis, Tunisia.

Received: 1 July 2020 Accepted: 28 January 2021

Published online: 03 February 2021

References

- Sohrabi C, Alsafi Z, O'Neill N, Khan M, Kerwan A, Al-Jabir A, et al. World health organization declares global emergency: a review of the 2019 novel coronavirus (COVID-19). *Int J Surg*. 2020;76:71–6.
- Organization WH. WHO announces COVID-19 outbreak a pandemic 2020 [2020 May 13]. Available from: <http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who-announces-covid-19-outbreak-a-pandemic>.
- Coronavirus Outbreak [15 May 2020]. Available from: <https://www.worldometers.info/coronavirus/>.
- Ouchetto O, Drissi Bourhanbour A, Boumhamdi M. Effectiveness of containment measures to control the spread of COVID-19 in North Africa. *Disaster Med Public Health Prep*. 2020. <https://doi.org/10.1017/dmp.2020.314>.
- COVID-19-dernier-bulletin.pdf [2020 May 14]. Available from: <http://www.onmne.tn/fr/images/COVID-19-dernier-bulletin.pdf>.
- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet*. 2020;395(10223):507–13.
- Young BE, Ong SWX, Kalimuddin S, Low JG, Tan SY, Loh J, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *Jama*. 2020;323(15):1488–94.
- Tian S, Hu N, Lou J, Chen K, Kang X, Xiang Z, et al. Characteristics of COVID-19 infection in Beijing. *J Infect*. 2020;80(4):401–6.
- Wong HYF, Lam HYS, Fong AH, Leung ST, Chin TW, Lo CSY, et al. Frequency and distribution of chest radiographic findings in patients positive for COVID-19. *Radiology*. 2020;296(2):E72–E8.
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *Jama*. 2020;323(11):1061–9.
- Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020;8(5):475–81.
- Wölfel RCV, Guggemos W, Seilmaier M, Zange S, Müller MA, et al. virological assessment of hospitalized patients with COVID-2019. *Nature*. 2020; 581(7809):465–9.

13. Organization H. Interim Guidance 2020 [3rd Apr 2020]. Available from: <https://apps.who.int/iris/bitstream/handle/10665/331329/WHO-COVID-19-laboratory-2020.4-eng.pdf>.
14. Choi WS, Kang CI, Kim Y, Choi JP, Joh JS, Shin HS, et al. Clinical presentation and outcomes of Middle East respiratory syndrome in the Republic of Korea. *Infect Chemother*. 2016;48(2):118–26.
15. Singhal T. A review of coronavirus Disease-2019 (COVID-19). *Indian J Pediatr*. 2020;87(4):281–6.
16. Zhao X, Zhang B, Li P, Ma C, Gu J, Hou P, et al. Incidence, clinical characteristics and prognostic factor of patients with COVID-19: a systematic review and meta-analysis. Preprint at <https://www.medrxiv.org/content/10.1101/2020.03.17.20037572v1>.
17. Guan WJ, Liang WH, Zhao Y, Liang HR, Chen ZS, Li YM, et al. Comorbidity and its impact on 1590 patients with COVID-19 in China: a nationwide analysis. *Eur Respir J*. 2020;55(5):2000547.
18. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72 314 cases from the Chinese Center for Disease Control and Prevention. *Jama*. 2020;323(13):1239–42.
19. Xu XW, Wu XX, Jiang XG, Xu KJ, Ying LJ, Ma CL, et al. Clinical findings in a group of patients infected with the 2019 novel coronavirus (SARS-CoV-2) outside of Wuhan, China: retrospective case series. *BMJ*. 2020;368:m606.
20. Andrea Giacomelli LP, Conti F, Bernacchia D, Siano M, Oreni L, et al. Self-reported olfactory and taste disorders in patients with severe acute respiratory coronavirus 2 infection: a cross-sectional study. *Clin Infect Dis*. 2020. <https://doi.org/10.1093/cid/ciaa330>.
21. Lee Y, Min P, Lee S, Kim SW. Prevalence and duration of acute loss of smell or taste in COVID-19 patients. *J Korean Med Sci*. 2020;35(18):e174.
22. Lechien JR, Chiesa-Estomba CM, De Siaty DR, Horoi M, Le Bon SD, Rodriguez A, et al. Olfactory and gustatory dysfunctions as a clinical presentation of mild-to-moderate forms of the coronavirus disease (COVID-19): a multicenter European study. *European archives of oto-rhino-laryngology: official journal of the European Federation of Oto-Rhino-Laryngological Societies*. 2020;277(8):2251–61.
23. Position Statement from the National Centre for Infectious Diseases and the Chapter of Infectious Disease Physicians, Academy of Medicine, Singapore: Period of Infectivity to Inform Strategies for De-isolation for COVID-19 Patients.pdf [2020 June 11]. Available from: <https://scholarbank.nus.edu.sg/handle/10635/168938>.
24. Xiao AT, Tong YX, Zhang S. False-negative of RT-PCR and prolonged nucleic acid conversion in COVID-19: Rather than recurrence. *J Med Virol*. 2020; 92(10):1755–6.
25. Li Y, Yao L, Li J, Chen L, Song Y, Cai Z, et al. Stability issues of RT-PCR testing of SARS-CoV-2 for hospitalized patients clinically diagnosed with COVID-19. *J Med Virol*. 2020;92(7):903–8.
26. Bullard J, Dust K, Funk D, Strong JE, Alexander D, Garnett L, et al. Predicting infectious SARS-CoV-2 from diagnostic samples. Preprint at <https://europepmc.org/article/med/32442256>.
27. Organization WH. Clinical management of severe acute respiratory infection (SARI) when COVID-19 disease is suspected. 2020 13 March 2020. Report No.
28. Mollazehi M, Mollazehi M, Abdel-Salam A-SG. Modeling Survival Time to Recovery from COVID-19: A Case Study on Singapore. Preprint at <https://www.researchsquare.com/article/rs-18600/v2>.
29. Kaijin Xu YC, Jing Y, Ping Yi, Cheng D, Wenrui Wu, et al. Factors Associated With Prolonged Viral RNA Shedding in Patients With COVID-19. *Clin Infect Dis*. 2020;71:799–806.
30. Liu K, Chen Y, Lin R, Han K. Clinical features of COVID-19 in elderly patients: a comparison with young and middle-aged patients. *J Infect*. 2020;80:e14.
31. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054–62.
32. Wang L, He W, Yu X, Hu D, Bao M, Liu H, et al. Coronavirus Disease 2019 in elderly patients: characteristics and prognostic factors based on 4-week follow-up. *J Infect*. 2020;80(6):639–45.
33. Rachel E, Jordan PA, Cheng KK. Covid-19: risk factors for severe disease and death. *BMJ*. 2020;368:m1198.
34. (ECDC) TECfDpaC. Novel Coronavirus (SARS-COV-2): Discharge criteria for confirmed COVID-19-cases. ECDC Technical Report. 2020.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

