

CORRECTION

Open Access



Correction to: Clinical characteristics in blood stream infections caused by *Klebsiella pneumoniae*, *Klebsiella variicola*, and *Klebsiella quasipneumoniae*: a comparative study, Japan, 2014–2017

Kazuo Imai^{1,2}, Noriomi Ishibashi^{2,3}, Masahiro Kodana⁴, Norihito Tarumoto^{1,2*}, Jun Sakai^{1,2}, Toru Kawamura⁴, Shinichi Takeuchi⁴, Yoshitada Taji⁵, Yasuhiro Ebihara⁵, Kenji Ikebuchi⁴, Takashi Murakami^{2,6}, Takuya Maeda^{2,6}, Kotaro Mitsutake^{2,3} and Shigefumi Maesaki^{1,2}

Correction to: BMC Infectious Diseases (2019) 19:946
<https://doi.org/10.1186/s12879-019-4498-x>

Following publication of the original article [1], the original PDF-publication of this article contained watermarking which indicated 'uncorrected proofs' whilst it was the final version.

The original article has been [1] updated to remove the watermarking.

Author details

¹Department of Infectious Disease and Infection Control, Saitama Medical University, 38 Morohongo, Moroyama-machi, Iruma-gun, Saitama 350-0495, Japan. ²Center for Clinical Infectious Diseases and Research, Saitama Medical University, 38 Morohongo, Moroyama-machi, Iruma-gun, Saitama, Japan. ³Infectious Diseases and Infection Control, Saitama Medical University International Medical Center, 1-1397 Yamane, Hidaka, Saitama 350-1298, Japan. ⁴Clinical Laboratory Medicine, Saitama Medical University Hospital, 38 Morohongo, Moroyama-machi, Iruma-gun, Saitama, Japan. ⁵Department of Clinical Laboratory Medicine, Saitama Medical University International Medical Center, 1-1397 Yamane, Hidaka, Saitama 1298, Japan. ⁶Department of Microbiology,

Saitama Medical University, 38 Morohongo, Moroyama-machi, Iruma-gun, Saitama 350-0495, Japan.

Accepted: 14 April 2022
Published online: 04 May 2022

Reference

1. Imai K, Ishibashi N, Kodana M, Tarumoto N, Sakai J, Kawamura T, Takeuchi S, Taji Y, Ebihara Y, Ikebuchi K, Murakami T, Maeda T, Mitsutake K, Maesaki S. Clinical characteristics in blood stream infections caused by *Klebsiella pneumoniae*, *Klebsiella variicola*, and *Klebsiella quasipneumoniae*: a comparative study, Japan, 2014–2017. *BMC Infect Dis*. 2019;19:946. <https://doi.org/10.1186/s12879-019-4498-x>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1186/s12879-019-4498-x>.

*Correspondence: tarumoto@saitama-med.ac.jp

¹ Department of Infectious Disease and Infection Control, Saitama Medical University, 38 Morohongo, Moroyama-machi, Iruma-gun, Saitama 350-0495, Japan

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.