

RESEARCH

Open Access



Individual and household risk factors for Ebola disease among household contacts in Mubende and Kassanda districts, Uganda, 2022

Stella M. Migamba^{1*}, Denis-Luc Ardiét², Richard Migisha¹, Hildah T. Nansikombi¹, Brian Agaba¹, Helen Nelly Naiga¹, Mercy Wanyana¹, Jane Frances Zalwango¹, Immaculate Atuhaire¹, Peter Chris Kawungezi¹, Marie Goretti Zalwango¹, Brenda Simbwa¹, Daniel Kadobera¹, Alex R. Ario¹ and Julie R. Harris³

Abstract

Background In 2022, an Ebola disease outbreak caused by Sudan virus (SUDV) occurred in Uganda, primarily affecting Mubende and Kassanda districts. We determined risk factors for SUDV infection among household members (HHM) of cases.

Methods We conducted a case-control and retrospective cohort study in January 2023. Cases were RT-PCR-confirmed SUDV infection in residents of Mubende or Kassanda districts during the outbreak. Case-households housed a symptomatic, primary case-patient for ≥ 24 h and had ≥ 1 secondary case-patient with onset < 2 weeks after their last exposure to the primary case-patient. Control households housed a case-patient and other HHM but no secondary cases. A risk factor questionnaire was administered to the primary case-patient or another adult who lived at home while the primary case-patient was ill. We conducted a retrospective cohort study among case-household members and categorized their interactions with primary case-patients during their illnesses as none, minimal, indirect, and direct contact. We conducted logistic regression to explore associations between exposures and case-household status, and Poisson regression to identify risk factors for SUDV infection among HHM.

Results Case- and control-households had similar median sizes. Among 19 case-households and 51 control households, primary case-patient death (adjusted odds ratio [OR_{adj}]=7.6, 95% CI 1.4–41) and ≥ 2 household bedrooms (OR_{adj}=0.19, 95% CI 0.056–0.71) were associated with case-household status. In the cohort of 76 case-HHM, 44 (58%) were tested for SUDV < 2 weeks from their last contact with the primary case-patient; 29 (38%) were positive. Being aged ≥ 18 years (adjusted risk ratio [aRR_{adj}]=1.9, 95%CI: 1.01–3.7) and having direct or indirect contact with the primary case-patient (aRR_{adj}=3.2, 95%CI: 1.1–9.7) compared to minimal or no contact increased risk of Sudan virus disease (SVD). Access to a handwashing facility decreased risk (aRR_{adj}=0.52, 95%CI: 0.31–0.88).

*Correspondence:
Stella M. Migamba
smigamba@musph.ac.ug

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Conclusion Direct contact, particularly providing nursing care for and sharing sleeping space with SVD patients, increased infection risk among HHM. Risk assessments during contact tracing may provide evidence to justify closer monitoring of some HHM. Health messaging should highlight the risk of sharing sleeping spaces and providing nursing care for persons with Ebola disease symptoms and emphasize hand hygiene to aid early case identification and reduce transmission.

Keywords Sudan virus disease, Ebola disease, Sudan virus, Uganda

Background

Ebolaviruses have the potential to cause both small and large outbreaks. Human Ebola disease (EBOD) outbreaks typically occur after humans have contact with the body fluids or meat of infected non-human primates [1] or fruit bats [2]. Person-to-person transmission occurs through direct contact with body fluids (such as blood, saliva, urine, sweat, vomit, faeces, breast milk, semen, vaginal fluid) or tissues of an infected, symptomatic person or corpse [3]. Among household contacts, EBOD is transmitted largely through direct physical contact with a symptomatic patient [4–6]. Transmission through contaminated inanimate objects (fomites) occurs, but is less common [5, 6]. A meta-analysis of household secondary attack rates for EBOD showed that risk was highest for household members providing nursing care (48%), and lowest for household members without any direct contact (0.8%) [5].

The first and largest EBOD outbreak in Uganda, caused by Sudan virus (SUDV), occurred in 2000 and included 425 cases and 224 deaths; additional EBOD outbreaks in Uganda occurred in 2007, 2011, 2012, and 2019 [7]. On September 20, 2022, the Uganda Ministry of Health (MoH) declared an outbreak of Sudan virus disease (SVD) after a case was confirmed the previous day in a 26-year-old man living in Mubende District, Central Uganda [8, 9]. The response to the outbreak included immediate and intensive efforts to control the outbreak and stop transmission, including rapid identification, isolation, and treatment of cases, contact tracing and monitoring of contacts [10, 11]. On October 15, in response to the spread to other districts, the president of Uganda instituted a 21-day lockdown in Mubende and Kassanda districts, the epicentres of the outbreak [12].

By the end of the outbreak, there were 142 confirmed cases of SVD in nine districts in Uganda [13]. In total, 4,793 contacts had been listed and monitored [14]. Household and community transmission accounted for two-thirds of cases [15]. As persons who typically interact the most closely with EBOD patients before they reach health facilities, household members of infected persons are at high risk of exposure during an outbreak. As in previous outbreaks [3, 6], during the 2022 SUDV outbreak in Uganda, some households had multiple cases. Due to the nonspecific symptoms of early infection, appropriate precautions may not be taken by

household members. EBOD symptoms are often similar to those of other diseases such as malaria or diarrhoeal diseases. Knowing which individual or household characteristics are associated with the highest risk of household transmission can provide information on tailored community education in an outbreak-affected area, and may provide information on contacts who need to be monitored especially carefully. We determined risk factors for Sudan virus (SUDV) infection among household members of confirmed cases in Mubende and Kassanda districts during the 2022 outbreak.

Methods

Study setting

This study was conducted in Mubende and Kassanda districts in Central Uganda, where 80% of the SUDV cases in the 2022 outbreak were identified [15]. The mid-year population projections for 2021 were 582,900 for Mubende District and 319,900 for Kassanda District [16]. Subsistence farming is the most common occupation [17].

Study design

We conducted both a case-control study and a retrospective cohort study in January 2023.

Case-control study

The case-control study was designed to understand household factors that increased the odds of presumed household transmission. Case-households were homes that housed a symptomatic, confirmed case-patient for at least 24 h and had a secondary case-patient with onset < 2 weeks after their last exposure to the primary case-patient (presumed infected from the primary case-patient). Control households were homes that housed a symptomatic, confirmed case-patient for at least 24 h but did not have another person who developed an infection at home at any point. Both case- and control households had to have at least one household member (HHM) besides the primary case-patient.

We identified case and control households from the SUDV confirmed case line list and narrative notes. Eligible households for the case-control study were selected based on the following criteria: First, the primary case-patient (the first person to develop confirmed SUDV in the household) had to have spent at least 24 h at home

while ill before being evacuated to the Ebola Treatment Unit (ETU). Second, the household had to have ≥ 1 HHM other than the primary case-patient who also spent at least 24 h in the home while the primary case-patient was ill and thus had a risk of becoming infected. All primary and secondary case-patients were laboratory-confirmed cases. Laboratory testing for SUDV was done at the Uganda Virus Research Institute (UVRI) laboratory, which was the designated national reference laboratory for viral haemorrhagic fever testing. SUDV infection was detected using real-time polymerase chain reaction (PCR) test [8, 13, 15].

Cohort study

Using only the case-households from the case-control study, we conducted a retrospective cohort study to identify individual risk factors for SUDV infection among all household members of primary case-patients. The cohort comprised all consenting case-HHM. We excluded HHM who could not be reached due to logistical reasons (relocation or not available by phone), who died without any next-of-kin available to interview, and who had mental disorders.

Data collection and study variables

For the case-control study, we administered questionnaires [18] to either the primary case-patient, if alive, or another adult who lived in the household while the primary case-patient was ill. We collected data on the primary case-patient's socio-demographic characteristics, presence of symptoms, number of days the primary case-patient was at home with symptoms, household location (urban or rural), number of rooms and bedrooms in the household, isolation practices for the primary case-patient, whether or not the primary case-patient had a single dedicated caretaker at home, presence of handwashing facilities, and type of care HHM provided to the primary case-patient (interactions).

For the cohort study, we collected data by interviewing every consenting adult HHM who lived in a case-household while the primary case-patient was ill, using a standardized questionnaire. For minors, we interviewed guardians, and for HHM who died, we interviewed a proxy. The objectives of these interviews were to characterise the level of interaction between household members and the primary case-patient. We collected data on HHMs' socio-demographic characteristics, whether they had underlying conditions or not, which symptoms they developed, whether they were tested for SUDV, whether they suspected that the primary case-patient had SVD, ways in which the contact interacted with the primary case-patient after his or her onset, use of gloves, access to a handwashing facility with soap and water, access to

information on the provision of safe care, and the clinical outcome of the primary case-patient.

Interactions were grouped into mutually exclusive groups to compare exposure-outcome associations against a common reference group. Interactions of the contact with the primary case-patient were categorised as no contact, minimal contact, indirect contact, and direct contact. 'No contact' referred to having had no interaction at all with the primary case-patient. 'Minimal contact' referred to having sat with or talked to the primary case-patient in the same room, or having removed dishes after meals or rode on the same motorbike, but none of the higher-level interactions. 'Indirect contact' included having washed the primary case-patient's clothes, changed their beddings, or cleaned their room but none of the higher-level interactions. 'Direct contact' including having played with the primary case-patient or bathed, cleaned, carried, helped move around, fed or breastfed, shared dishes or utensils at meals, or shared a bed, or had sexual intercourse with the primary case-patient.

We considered every HHM who was tested for SUDV infection and received a positive result to have SVD and every other contact to be negative (including those who were not tested). None of the HHM who were not tested reported any symptoms.

Data analysis

Case-control study

We conducted logistic regression to explore possible associations between each exposure variable and case-household status. Odds ratios and their associated 95% confidence intervals were used as measures of effect size. Exposures with p -values < 0.2 were included in the multivariable model. Multivariable analysis was done to determine predictors of being a case household.

Cohort study

We fitted generalized linear models using Poisson regression analysis to identify risk factors for SUDV infection among household members to primary case-patients and adjusted for clustering at household level. We included variables as categorical fixed effects nested within fixed household identifiers and assumed a normal distribution of the random effects.

We computed risk ratios with 95% confidence intervals to determine associations between exposures and confirmed SUDV infection. Exposures with p values < 0.2 were evaluated in multivariable analysis after checking for collinearity of variables to determine factors independently associated with SUDV infection. Stata version 14 (StataCorp, CollegeTexas, USA) was used for statistical analysis.

Results

Characteristics of case- and control-households

During the outbreak, there were 84 households with ≥ 1 case-patient, of which 70 households were eligible for the case-control study (i.e., had at least one HHM besides the primary case-patient). Of these, 19 were case-households and 51 were control households. Twelve (63%) case-households and 38 (75%) control households had a primary case-patient aged ≥ 18 years. The primary case-patient in 17 (89%) case-households and 28 (55%) control households died. Case- and control households had similar median household sizes (6 HHM, IQR 4–9 for case-households and 6 HHM, IQR 4–8 for control households). Ten (67%) case-households and 23 (62%) control households had at least 6 HHM (Table 1). All households (100%) reported at least one kind of care interaction of HHM with the primary case-patient.

Factors associated with household SUDV transmission

In multivariable analysis, households in which the primary case-patient died (whether at home or in hospital) had nearly eight times higher odds of becoming case-households than those in which the primary case-patient recovered ($OR_{adj}=7.6$, 95% CI: 1.4–41). Households with ≥ 2 bedrooms had lower odds of being case-households than those that had only one bedroom ($OR_{adj}=0.19$, 95% CI: 0.056–0.71) (Table 1).

Cohort characteristics

From 19 case-households in the case-control study, we enrolled 76 of 108 total household members for the cohort study (Fig. 1).

Mean HHM age was 24 (± 17) years. Mean number of days from primary case-patient's reported onset date to the HHM's onset date was 8 (range, 1–20); median was 7 (IQR 4–10). A total of 44 (58%) HHM were tested for SUDV infection. Twenty-nine (38%) HHM overall who had illness onset ≤ 2 weeks from their last contact with the primary case-patient tested positive for SUDV infection (Table 2).

By individual interaction (not mutually exclusive), infection rates were highest among those who had sexual intercourse with the primary case-patient during his or her illness (Table 3).

Among the 76 household members, 13 (17%) had no contact with the primary case-patient in their household; none of these 13 became ill. Four (5%) had minimal contact, of whom two became ill. Three (4%) had indirect contact, and one became ill. Fifty-six (74%) had direct contact, and 26 became ill (Table 4).

Household members who had direct contact with the primary case-patient in their households had a three-fold higher risk of contracting SUDV infection than those who had either no, or minimal or indirect contact only.

Additionally, those who had direct and/or indirect contact had four times the risk of SUDV infection compared to those who had no contact and/or minimal contact (Table 5).

In multivariable analysis, being aged ≥ 18 years ($aRR_{adj}=1.9$, 95% CI: 1.01–3.7) and having had direct and/or indirect contact but not minimal contact with the primary case-patient ($aRR_{adj}=3.2$, 95% CI: 1.1–9.7) increased the risk of SUDV infection among household members. Access to a handwashing facility decreased the risk of SUDV infection ($aRR_{adj}=0.52$, 95% CI: 0.31–0.88) (Table 6).

Risk factors for SUDV infection among HH members

Discussion

This study found multiple factors to be associated with SUDV infection among household members of confirmed cases during the 2022 outbreak in Uganda. Having a case-patient die increased odds of a household having secondary SUDV cases. Households that had more than one bedroom had lower odds of having secondary infections than those that had only one bedroom. Higher levels of contact with the case-patient, especially nursing care, shared sleeping space, and sexual contact were associated with increased infection risk among household members. Being an adult household member and not having access to a handwashing facility increased risk of infection.

Nearly 90% of primary case-patients in case-households died, compared to slightly more than half of those in control households. This finding is similar to those from two studies in Sierra Leone where index patient death was a risk factor for household transmission [19, 20]. The association between primary case-patient death and secondary cases among household members may be due to the increased infectiousness during advanced disease [6] as a result of an increase in viral load [21]. Death among patients with EBOD has been associated with delays to care [22], which may be correlated with a longer time at home and increased time to expose household members. Interestingly, in our study, we did not identify differences between case-households and control households in the amount of time the primary case-patient spent at home while ill, which may suggest that the primary case-patients in case-households faced more rapid disease progression than those in control households. However, we lacked sufficient clinical data to assess this possibility.

Households that had more than one bedroom had lower odds of having secondary infections than those that had only one bedroom. However, the odds of SUDV infection did not differ significantly by the total number of rooms in the household or by the ratio of household

Table 1 Characteristics of case and control households during the Ebola outbreak in Mubende and Kassanda districts, Uganda, 2022

Variable (n = 70)	Case HH n (%)	Control HH n (%)	cOR (95% CI)	aOR (95% CI)
Primary case-patient sex				
Male	12 (63)	27 (53)	1	
Female	7 (37)	24 (47)	0.66 (0.22–1.9)	
Age of primary case-patient (yrs)				
< 18	7 (37)	13 (25)	1	1
≥ 18	12 (63)	38 (75)	0.59 (0.19–1.8)	0.67 (0.18–2.5)
Clinical outcome of primary case-patient				
Recovered	2 (11)	23 (45)	1	1
Died	17 (89)	28 (55)	6.9 (1.5–33)	7.6 (1.4–41)
District				
Mubende	12 (63)	36 (71)	1	
Kassanda	7 (37)	15 (29)	1.4 (0.46–4.2)	
HH location				
Urban	6 (32)	19 (37)	1	
Rural	13 (68)	32 (63)	1.3 (0.42–3.9)	
HH bedrooms				
1	10 (53)	11 (22)	1	1
≥ 2	9 (47)	40 (78)	0.25 (0.081–0.76)	0.19 (0.056–0.71)
HH rooms				
1	4 (21)	6 (12)	1	
≥ 2	15 (79)	45 (88)	0.50 (0.12–2.01)	
Number of HHM*				
2–5	9 (47)	14 (38)	1	
≥ 6	10 (53)	23 (62)	0.67 (0.22–2.07)	
Ratio HHM: bedrooms				
1:1–4:1	13 (68)	31 (84)	1	
> 4:1	6 (32)	6 (16)	2.4 (0.65–8.8)	
Ratio HHM: rooms				
1:1–2:1	12 (63)	20 (54)	1	
3:1–9:1	7 (37)	17 (46)	0.69 (0.22–2.1)	
Household had one dedicated caretaker for primary case-patient				
No	10 (53)	24 (47)	1	1
Yes	9 (47)	27 (53)	0.80 (0.28–2.3)	0.81 (0.24–2.7)
Household had piped water for handwashing				
No	17 (89)	46 (90)	1	
Yes	2 (11)	5 (10)	1.1 (0.19–6.1)	
Household had electricity				
No	5 (26)	12 (24)	1	
Yes	14 (74)	39 (76)	0.86 (0.26–2.9)	
Primary case-patient had an underlying condition				
No	16 (84)	38 (75)	1	
Yes	3 (16)	13 (25)	0.55 (0.14–2.2)	
Days primary case-patient was ill at home before evacuation				
1	2 (10)	5 (10)	1	1
2–4	7 (37)	18 (35)	0.97 (0.15–6.2)	1.5 (0.18–13)
≥ 5	10 (53)	28 (55)	0.89 (0.15–5.4)	1.6 (0.21–12)
Primary case-patient stayed isolated at home during illness				
No	18 (95)	45 (88)	1	
Yes	1 (5)	6 (12)	0.42 (0.047–3.7)	

*Of the 70 households, data on household size were only available for 56, including 19 case-households and 37 control households; HH: Household; HHM: Household member; cOR: Crude odds ratio; aOR: Adjusted odds ratio




Fig. 1 Persons enrolled in the cohort study of risk factors for SUDV infection, Uganda, 2022

members to either total rooms or bedrooms. This suggests that while crowding itself might not have increased risk in our study, shared sleeping space was a specific risk. In support of this finding, assessment of individual (non-mutually exclusive) interactions showed that infection rates were highest among those who had sex with the primary case-patient (67%) followed by those who shared a bed with the primary case-patient (59%). Infection during sexual intercourse may have occurred due to the requisite physical intimacy of the act, or possibly through sexual transmission. Although sexual transmission has only been documented from survivors [23], the virus is known to be present in body fluids including semen and vaginal fluids, and this is a possible mechanism of transmission. However, it is likely that persons having sexual intercourse with a patient also had other (nonsexual) exposures to the patient, possibly later in the illness, that could have put them at risk. In addition, only three of the 37 household members sharing a bed with the primary case-patient reported sexual intercourse with the primary case-patient, and infection rates were similar between these two groups. This suggests that simple proximity during sleeping may suffice to transmit infection, either due to physical contact or fomite contact.

In agreement with other studies demonstrating that close physical contact — and specifically nursing care — is a strong risk factor for infection [5, 24], we found that bathing or cleaning a patient, carrying, helping to move around, playing with, feeding or breastfeeding a patient, and sharing utensils at meals with the primary case-patient all increased risk. In the absence of direct contact, risk of infection reduced greatly, and no household members without any contact developed infection; this is

consistent with other studies [5, 24]. While fomites can serve as a source of ebolavirus infection, this study adds to the body of evidence that this is a less common infection pathway than direct contact [5, 6].

In the cohort study, adult household members (18 years or older) were more likely to contract SUDV infection than those younger than 18 years. This finding is consistent with those from two studies that revealed that children are usually less affected than adults during Ebola outbreaks [3, 5, 25]. In contrast, Fang et al. [26] found that children had higher odds of household infection than adults, although this finding was not statistically significant. A possible explanation for children being less affected could be their limited ability to provide nursing care to ill family members, reducing their chances of a high-risk exposure. However, children — especially those <5 years of age — have a higher case-fatality rate than adults from EBOD [27–29], and measures should be instituted to protect them even when they do not provide direct care to a patient.

In our cohort study, access to a handwashing facility reduced risk of infection among household members by half. This is not surprising; ebolavirus infections are primarily transmitted through contact with contaminated body fluids [4–6]. Handwashing can reduce risk by supporting both the patient's own hygiene as well as the hygiene of his or her caretakers. Inclusion of messaging that emphasizes hand hygiene in communities affected by EBOD continues to be important.

Our study had several limitations. First, due to varied incubation periods, it is possible that some household members within a single household may have been infected by a common source external to the household and had different onsets. For example, siblings caring for an ill mother living outside the home might have both acquired infection from the mother at a similar time or perhaps sequentially. The median reported serial interval between the primary and secondary cases in our study was 7 days, four days shorter than the reported serial interval for the outbreak overall [30], suggesting that this limitation may have applied to at least some of our secondary cases. This would have made the associations between primary and secondary cases in case-households appear stronger than they really are. Second, in some households (particularly when a case-patient had died), a proxy was interviewed. The proxy might not have had accurate knowledge of interactions that may have occurred between the primary case-patient and household members, or of specific dates. Recall bias might also have resulted in under-reporting of some interactions. Finally, some sensitive interactions, such as sexual contact, may have been under-reported and thus the risk they posed may not have fully been assessed.

Table 2 Characteristics of household members (HHM) (n = 76) of primary SVD cases in Mubende and Kassanda districts, Uganda, 2022

Variable	Frequency	%
Age		
< 18	33	43
≥ 18	43	57
Mean age (± SD)	24 (± 17)	
Sex		
Male	30	39
Female	46	61
Relationship of HHM to primary case-patient		
Daughter/son	15	20
Mother/ father	18	24
Sibling	18	24
Husband/ wife	8	11
Aunt/ uncle	4	5
Grandchild	7	9
Other	6	8
Occupation of HHM		
Child or student	36	47
Other professions	40	53
Highest education attained		
None and primary	67	88
Secondary and above	9	12
HHM developed signs/symptoms of SVD (self-reported)		
Yes	32	42
No	44	58
Days from primary case-patient illness onset to HHM onset		
Median (IQR)	7 (4-10)	
HHM tested for SUDV		
Yes	44	58
No	32	42
SUDV test result (among all household members)		
Positive	29	38
Negative	47	62
HHM had an underlying condition		
Yes	8	11
No	68	89
HHM interacted with primary case-patient during illness		
Yes	63	83
No	13	17
Household suspected primary case-patient had SVD		
Yes	3	4
No	73	96
HHM had gloves		
Yes	2	3
No	74	97
HHM had access to handwashing station with soap		
Yes	60	79
No	16	21
Frequency of handwashing^a		
Less than half of the time	47	78
More than half the time/ all the time	13	22
HHM tried to keep distance from primary case-patient²		
Yes	5	7
No	64	93

Table 2 (continued)

Variable	Frequency	%
HHM knew how to interact with primary case-patient safely²		
Yes	3	4
No	66	96
HHM was given information on caring for primary case-patient safely^b		
Yes	4	6
No	65	94

^an=60; ^bn=69; SD: Standard deviation; IQR: Interquartile range; HH: Household; HHM: Household member; SVD: Sudan virus disease; SUDV: Sudan virus

Table 3 Interactions between household members and primary case-patients in Mubende and Kassanda districts, Uganda, and subsequent positive test proportion, 2022 (n = 63)

Interaction	Total, n	SUDV +, n	(%)
Had sexual intercourse with primary case-patient	3	2	(67)
Shared bed with primary case-patient	37	22	(59)
Carried/ held primary case-patient	27	16	(59)
Bathed/ cleaned primary case-patient	31	18	(58)
Helped primary case-patient move around	21	12	(57)
Fed primary case-patient	27	15	(56)
Cleaned primary case-patient's room	20	11	(55)
Changed primary case-patient's beddings	35	19	(54)
Washed primary case-patient's clothes	36	19	(53)
Breastfed primary case-patient	2	1	(50)
Removing primary case-patient's dishes	47	23	(49)
Played with primary case-patient	13	6	(46)
Shared utensils with case-patient at meals	42	18	(43)
Rode on boda with primary case-patient	7	3	(43)
Sat with primary case-patient in same room	27	10	(37)
Exchanged money with primary case-patient	7	0	(0)

Conclusion

Direct contact, and particularly sharing sleeping space or providing nursing care to an EBOD patient, increased risk of infection among household members. Health messaging during EBOD outbreaks should emphasize the risk of both sleeping next to and providing nursing care for persons with symptoms consistent with EBOD, even early EBOD, until testing can rule out infection. However, we note that in the early stages of SUDV outbreak, direct contact of household members with primary case-patients may be inevitable. Often, before a diagnosis is made, unsuspecting household members have already been exposed to infectious primary case-patients. Implementation of risk assessments for household contacts may provide important data to justify closer monitoring of those considered to be at especially high-risk during contact tracing activities.

Table 4 SUDV positivity by level of contact among household members of primary case-patients in Mubende and Kassanda districts, Uganda, 2022 (n = 76). Interaction levels represent the maximum degree of interaction between the household member and primary case-patient

Interaction category	Total (col %) n (%)	SUDV + (row %) n (%)
No contact	13 (17)	0 (0)
Minimal contact	4 (5)	2 (50)
Indirect contact	3 (4)	1 (33)
Direct contact	56 (74)	26 (46)

Table 5 Grouped interactions of household members of primary case-patients in Mubende and Kassanda districts, Uganda, 2022

Exposure (n = 76)	n	SUDV +	uRR* (95% CI)	P value
Any direct contact vs. (no contact, minimal contact, or indirect contact only)				
No, minimal, or indirect contact	20	3	1	
Direct contact	56	26	3.1 (1.1–9.1)	0.042
Any direct or indirect contact vs. (no contact or minimal contact only)				
No contact or minimal contact	17	2	1	
Direct and indirect contact	59	27	3.9 (1.02-15)	0.047

*uRR – unadjusted risk ratio; SUDV: Sudan virus

Table 6 Risk factors for SVD among household contacts to primary case-patients in Mubende and Kassanda districts, Uganda, 2022

Exposure (n = 76)	n	SUDV +	uRR (95% CI)	aRR (95% CI)	P value
Age					
< 18	42	22	1	1	
≥ 18	34	8	2.1 (1.08–4.2)	1.9 (1.01–3.7)	0.05
Access to handwashing facility					
No	16	10	1	1	
Yes	60	19	0.51 (0.29–0.86)	0.52 (0.31–0.88)	0.013
Sex					
Male	30	11	1	1	
Female	46	18	1.07 (0.59–1.9)	1.1 (0.63–1.9)	0.69
Highest education					
None and primary	67	26	1	1	
Secondary and above	9	3	0.86 (0.33–2.3)	0.84 (0.35–2.0)	0.70
Primary case-patient's outcome					
Died	24	12	1	1	
Recovered	52	17	0.65 (0.37–1.1)	0.8 (0.44–1.4)	0.38
Direct or indirect contact vs. (no contact or minimum contact) *					
No contact or minimal contact	17	2	1	1	
Direct or indirect contact	59	27	3.9 (1.02–15)	3.2 (1.1–9.7)	0.004

*Representing the maximum level of contact a household member had with a case-patient

uRR: unadjusted risk ratio; aRR: adjusted risk ratio; SUDV: Sudan virus

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12879-024-09439-1>.

Supplementary Material 1

Supplementary Material 2

Acknowledgements

We acknowledge the case investigation teams that collected narratives on case-patients as well as the local council chairpersons and community health workers of Mubende and Kassanda districts who guided in locating households.

Author contributions

SMM, DA and JRH conceptualised the study and designed the analysis plan. PCK, RM, AB, NN, MW, JFZ, IA, GMZ, and BS reviewed the data collection tools. SMM and JRH analysed the data. HTN advised on the analysis plan. RM, JRH, DK, and ARA contributed substantially to revisions of the manuscript. All authors were part of the outbreak investigation and response.

Funding

This study was supported by the President's Emergency Plan for AIDS Relief (PEPFAR) through US Centre for Disease Control and Prevention Cooperative Agreement number GH001353–01 through Makerere University School of Public Health to the Uganda Public Health Fellowship Program, Ministry of Health. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the U.S. Centers for Disease Control and Prevention.

Data availability

The data upon which our findings are based belongs to the Uganda Ministry of Health and cannot be shared publicly for confidentiality reasons. However, it can be made available by the corresponding author with permission from the Ministry of Health Uganda, Division of Health Information and Uganda Public Health Fellowship Program.

Declarations

Ethics approval and consent to participate

This outbreak investigation was in response to a public health emergency. The Ministry of Health gave a directive to investigate this outbreak and the office of the Center for Global Health, US Center for Disease Control and Prevention determined that this activity was non-research and that its primary intent was for public health practice or disease control. This analysis of the dynamics of transmission of SUDV to household members from primary case-patients was done as part of integrated outbreak analytics (IOA). This activity was reviewed by CDC and was conducted consistent with applicable federal law and CDC policy. § See e.g., 45 C.F.R. part 46, 21 C.F.R. part 56; 42 U.S.C. § 241(d); 5 U.S.C. § 552a; 44 U.S.C. § 3501 et seq.

All SUDV testing was done for diagnostic purposes as part of the public health emergency response to the outbreak in Uganda, and thus was not subject to institutional review board clearance. Test results were released by the Uganda Virus Research Institute (UVRI) laboratory to the relevant public health authorities according to protocol. We included households and individuals who consented to the interviews. The authors sought permission to conduct the study from district health authorities and local government leaders of Mubende and Kassanda districts.

Verbal informed consent was sought from study participants who were at least 18 years old or their parents/guardians for those below 18 years. The authors also sought assent from participants below the age of 18 years. For those who had died, informed consent was obtained from their next-of-kin. The interviewers took the participants/guardians through the informed consent section and obtained consent before proceeding with interviews. The choice of verbal over written consent was made so as to minimise the possible risk of Ebola transmission. Each study participant was assigned a unique identifier to protect their confidentiality. All methods were conducted consistent with relevant guidelines and regulations.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

Author details

¹Uganda Public Health Fellowship Program, Uganda National Institute of Public Health, Kampala, Uganda

²Department of Epidemiology and Training, Epicentre, Paris, France

³Division of Global Health Protection, U.S. Centers for Disease Control and Prevention, Kampala, Uganda

Received: 23 February 2024 / Accepted: 27 May 2024

Published online: 30 May 2024

References

- Geisbert TW, Strong JE, Feldmann H. Considerations in the use of nonhuman primate models of Ebola virus and Marburg virus infection. *J Infect Dis*. 2015;212(suppl2):S91–7.
- Feldmann H, Geisbert TW. Ebola haemorrhagic fever. *Lancet*. 2011;377(9768):849–62.
- Borchert M, Mutyaba I, Van Kerkhove MD, Lutwama J, Luwaga H, Bisoborwa G, et al. Ebola haemorrhagic fever outbreak in Masindi District, Uganda: outbreak description and lessons learned. *BMC Infect Dis*. 2011;11(1):1–17.
- Bower H, Smout E, Bangura MS, Kamara O, Turay C, Johnson S et al. Deaths, late deaths, and role of infecting dose in Ebola virus disease in Sierra Leone: retrospective cohort study. *BMJ*. 2016;353.
- Dean NE, Halloran ME, Yang Y, Longini IM. Transmissibility and pathogenicity of Ebola virus: a systematic review and meta-analysis of household secondary attack rate and asymptomatic infection. *Clin Infect Dis*. 2016;62(10):1277–86.
- Francesconi P, Yoti Z, Declich S, Onek PA, Fabiani M, Olango J, et al. Ebola hemorrhagic fever transmission and risk factors of contacts. *Uganda Emerg Infect Dis*. 2003;9(11):1430.
- CDC. History of Ebola Virus Disease (EVD). Outbreaks 2022 [cited 2022 5 December]. <https://www.cdc.gov/vhf/ebola/history/chronology.html>.
- Kiggundu T, Ario AR, Kadobera D, Kwesiga B, Migisha R, Makumbi I, et al. Notes from the field: outbreak of Ebola virus disease caused by Sudan ebolavirus—Uganda, August–October 2022. *Morb Mortal Wkly Rep*. 2022;71(45):1457.
- WHO. Uganda declares Ebola Virus Disease outbreak: WHO Regional Office for Africa. 2022. <https://www.afro.who.int/countries/uganda/news/uganda-declares-ebola-virus-disease-outbreak>.
- Rai A, Hamiidah N, Abbass M, Masunga DS, Shoumar H, Kantawala BI et al. Ebola Virus Disease in Uganda: a global emergency call. *Annals Med Surg*. 2022;104825.
- WHO. WHO bolsters Ebola disease outbreak response in Uganda. 2022. <https://www.afro.who.int/countries/uganda/news/who-bolsters-ebola-disease-outbreak-response-uganda>.
- European Centre for Disease Prevention and Control. Ebola disease outbreak caused by Sudan ebolavirus in Uganda. 2022.
- Musoke P, Bongomin F. Sudan virus disease outbreak in Uganda in 2022: the case of patient zero. *Int J Infect Dis*. 2023;128:318–20.
- WHO. Ebola Virus Disease in Uganda Sitrep-93 2023. <https://www.afro.who.int/countries/uganda/publication/ebola-virus-disease-uganda-sitrep-93>.
- Kabami Z, Riolexus Ario A, Harris J, Ninsiima M, Ahirirwe SR, Aceng JR et al. Epidemiology of Cases of Sudan Virus Disease in Uganda, August–November, 2022.
- Uganda Bureau of Statistics. Population and, Censuses UBOS. 2024 [cited 2024 8 January]. <https://www.ubos.org/explore-statistics/20/>.
- Uganda Bureau of Statistics. The National Population and Housing Census 2014- Area Specific Profile Series. Kampala; 2014.
- Stella M, Migamba D-L, Ardiet R, Migisha, Hildah T, Nansikombi B, Agaba JR, Harris et al. Questionnaire_Individual and household risk factors for Ebola disease among household contacts in Mubende and Kassanda districts. Uganda2023.
- Glynn JR, Bower H, Johnson S, Turay C, Sesay D, Mansaray SH, et al. Variability in intrahousehold transmission of Ebola virus, and estimation of the household secondary attack rate. *J Infect Dis*. 2018;217(2):232–7.
- Reichler MR, Bruden D, Thomas H, Erickson BR, Knust B, Duffy N, et al. Ebola patient virus cycle threshold and risk of household transmission of Ebola virus. *J Infect Dis*. 2020;221(5):707–14.
- Towner JS, Rollin PE, Bausch DG, Sanchez A, Cray SM, Vincent M, et al. Rapid diagnosis of Ebola hemorrhagic fever by reverse transcription-PCR in an outbreak setting and assessment of patient viral load as a predictor of outcome. *J Virol*. 2004;78(8):4330–41.
- Qin E, Bi J, Zhao M, Wang Y, Guo T, Yan T, et al. Clinical features of patients with Ebola virus disease in Sierra Leone. *Clin Infect Dis*. 2015;61(4):491–5.
- Jacob ST, Crozier I, Fischer WA, Hewlett A, Kraft CS, Vega M-AL, et al. Ebola virus disease. *Nat Reviews Disease Primers*. 2020;6(1):13.
- Brainard J, Hooper L, Pond K, Edmunds K, Hunter PR. Risk factors for transmission of Ebola or Marburg virus disease: a systematic review and meta-analysis. *Int J Epidemiol*. 2016;45(1):102–16.
- Kourtis AP, Appelgren K, Chevalier MS, McElroy A. Ebola virus disease: focus on children. *Pediatr Infect Dis J*. 2015;34(8):893.
- Fang L-Q, Yang Y, Jiang J-F, Yao H-W, Kargbo D, Li X-L et al. Transmission dynamics of Ebola virus disease and intervention effectiveness in Sierra Leone. *Proceedings of the National Academy of Sciences*. 2016;113(16):4488–93.
- Agua-Agum J, Ariyaratna A, Blake IM, Cori A, Donnelly CA, Dorigatti I, et al. Ebola virus disease among children in West Africa. *N Engl J Med*. 2015;372(13):1274–7.
- Shah T, Greig J, van der Plas LM, Achar J, Caleo G, Squire JS, et al. Inpatient signs and symptoms and factors associated with death in children aged 5 years and younger admitted to two Ebola management centres in Sierra Leone, 2014: a retrospective cohort study. *Lancet Global Health*. 2016;4(7):e495–501.
- Smit MA, Michelow IC, Glavis-Bloom J, Wolfman V, Levine AC. Characteristics and outcomes of pediatric patients with Ebola virus disease admitted to treatment units in Liberia and Sierra Leone: a retrospective cohort study. *Clin Infect Dis*. 2017;64(3):243–9.
- Marziano V, Guzzetta G, Longini I, Merler S. Estimates of serial interval and Reproduction Number of Sudan Virus, Uganda, August–November 2022. *Emerg Infect Dis*. 2023;29(7):1429.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.