

POSTER PRESENTATION

Open Access

# Integrations as surrogate marker for multi drug resistance in gram negative bacilli

Gopalakrishnan Kaushik<sup>1\*</sup>, K Arunkumar<sup>2</sup>, G Sivakumar<sup>2</sup>, R Rajasomasundharam<sup>2</sup>, Padma Krishnan<sup>1</sup>

From First International Science Symposium on HIV and Infectious Diseases (HIV SCIENCE 2012) Chennai, India. 20-22 January 2012

## Background

Multi drug resistance (MDR) in gram negative bacilli (GNB) is an increasing global problem, which is often due to the acquisition of resistance genes from a shared pool. In MDR isolates, **integrations**-resistant gene cassettes, together with its accessory genetic elements (integrase) may be found in complex conglomerations on plasmids or on the chromosome and are widely acknowledged for their role in conferring MDR phenotype. Hence, the present study was done to evaluate the usefulness of integrase gene PCR as a surrogate marker for MDR in GNB.

## Methods

A total of 59 gram negative isolates collected between May 2010 and January 2011 from a tertiary hospital were included for this study. MDR phenotype of the isolates was established by the Kirby Bauer disc diffusion susceptibility test. Integrase gene PCR for class I and II integrations was performed.

## Results

Of the 59 isolates tested, *Pseudomonas* and *E. coli* were the predominant and 32 were MDR GNB. A total of 39 isolates were positive for either class I or class II integrations. Out of the 13 antibiotics tested, isolates carrying either class I or II integrations were resistant to 6 or more antibiotics, whereas isolates that were negative for carriage of integrations were resistant to 2 antibiotics or below.

## Conclusion

The study clearly indicates that integrations are widely prevalent in MDR GNB and further substantiates the

usefulness of integrase gene PCR as a surrogate marker for MDR in GNB.

## Author details

<sup>1</sup>Dept of Microbiology, Dr ALM PG Institute of Basic Medical Sciences, University of Madras, Chennai, India. <sup>2</sup>Department of General Surgery, Madras Medical College, Chennai, India.

Published: 4 May 2012

doi:10.1186/1471-2334-12-S1-P72

**Cite this article as:** Kaushik et al.: Integrations as surrogate marker for multi drug resistance in gram negative bacilli. *BMC Infectious Diseases* 2012 12(Suppl 1):P72.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [kaushikgangul@gmail.com](mailto:kaushikgangul@gmail.com)

<sup>1</sup>Dept of Microbiology, Dr ALM PG Institute of Basic Medical Sciences, University of Madras, Chennai, India

Full list of author information is available at the end of the article