

# **POSTER PRESENTATION**

**Open Access** 

# Rapid speciation of mycobacteria with simultaneous detection of MDR tB

Barnali Kakati<sup>\*</sup>, Rajiv Kumar Agarwal, Biswaroop Chatterjee, Aarti Kotwal, Pronoti Sarkar, Sudhir Singh, Victor Masih, Moirangthem Ajit Singh

From 2nd International Science Symposium on HIV and Infectious Diseases (HIV SCIENCE 2014) Chennai, India. 30 January - 1 February 2014

### **Background**

Multidrug resistance (MDR) TB is on the rise globally and is associated with significant morbidity and mortality. Conventional culture based drug susceptibility testing (DST) for TB drugs is time consuming, leading to diagnostic delays contributing to higher TB morbidity and mortality and exacerbation of ongoing transmission. Moreover most of the NTM(Non tuberculous mycobacteria) that are resistant to common anti tB drugs, can be falsely labeled as MDR tB if not identified correctly. This study was aimed to identify *Mycobacterium* species and to determine the prevalence of multidrug resistant tuberculosis (MDR tB) by line probe assay.

## Methods

This study was done on 76 clinical isolates collected from TB patients who were on Anti tuberculous drug treatment since last six months but without any clinical improvement. Line probe assay was performed for detection, identification of species and drug susceptibility status.

#### **Results**

Line probe assay detected 49 isolates (96%) of M. tuberculosis, 1 isolate (1.96%) of M. avium and M. intracellulare each. Out of 49 isolates of M. tuberculosis 9(18.4%) were Rifampicin resistant and 11(22.4%) INH resistant but what was alarming was that 9 isolates (18.4%) were MDR.

#### Conclusion

The majority of multidrug resistant TB cases are due to ongoing transmission of multidrug resistant strains. This is most likely the result of diagnostic delay, thereby

\* Correspondence: drbarnali05@rediffmail.com
Department of Microbiology, Himalayan Institute of Medical Sciences,
Dehradun-248140. India

emphasizing the need for rapid diagnostics and line probe assay can be used for the rapid identification of *Mycobacterium* species and the determination of susceptibility to drugs, which may accelerate the time to effective MDR tB treatment.

Published: 27 May 2014

doi:10.1186/1471-2334-14-S3-P43

Cite this article as: Kakati *et al.*: Rapid speciation of mycobacteria with simultaneous detection of MDR tB. *BMC Infectious Diseases* 2014 14(Suppl 3):P43.

# Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



