

POSTER PRESENTATION

Open Access

Emerging MDR producing ESBL among Diarrhoegenic *Esherichia coli* from paediatric patients

Ashita A Pinto*, Yaser Siahbalaei, Arif Ali

From 2nd International Science Symposium on HIV and Infectious Diseases (HIV SCIENCE 2014) Chennai, India. 30 January - 1 February 2014

Background

The beta lactam antibiotics are predominant therapeutic agents against a number of infectious diseases. The production of the enzyme beta lactamase by microbes which hydrolyses the beta lactam antibiotics is an important mechanism of resistance. This study aimed to determine prevalence, phenotypic patterns, and ESBL-production status both phenotypic and genotypic for the CTX –M type ESBL in *Escherichia coli* (*E.coli*) isolated from Diarrhoeic children.

Methods

Two hundred thirty non repetitive stool samples were collected from 2 hospitals, one private and other government hospital in Delhi, NCR. *E.coli* was confirmed by *16S* rRNA analyses. Antibiotic susceptibility and phenotypic ESBL production were studied by disc diffusion and double disk synergy tests according to Clinical Laboratory Standards Institute guidelines. PCR was performed for the molecular detection of the CTX-M type ESBL.

Results

Out of 230, 215 were confirmed to be *E.coli*. Phenotypically 72.5% (156) showed the presence of ESBL of which 75% (79) in private hospital samples and 70% (77) in government hospital samples. PCR based molecular detection confirmed the presence of the *CTX-M* gene among 65.6% (141) of the samples.

Conclusion

The high trends in the ESBL presence are alarming and urge means to cope with such ESBL *E.coli* strains.

* Correspondence: apintodelhi@gmail.com Gene Expression Laboratory, Department of Biosciences, Jamia Millia Islamia, New Delhi. India It being a commensal raises concern of the resistance gene transfer from pathogenic bacteria, especially in children due to common diarrhoeal episodes among them. The paediatric population is at a higher risk in case of multidrug resistance due to further limitation in therapeutic options and constrains in prescription of harsher antibiotics.

Published: 27 May 2014

doi:10.1186/1471-2334-14-S3-P70

Cite this article as: Pinto et al.: Emerging MDR producing ESBL among Diarrhoegenic Esherichia coli from paediatric patients. BMC Infectious Diseases 2014 14(Suppl 3):P70.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at www.biomedcentral.com/submit



