

RESEARCH ARTICLE

Open Access



# Exploring *Leishmania infantum* cathepsin as a new molecular marker for phylogenetic relationships and visceral leishmaniasis diagnosis

Ryan Emiliano da Silva<sup>1</sup>, Bruna Matarucco Sampaio<sup>1</sup>, Renata Tonhosolo<sup>2</sup>, Andrea Pereira da Costa<sup>3</sup>, Luiz Eduardo da Silva Costa<sup>1</sup>, Fernanda Ap. Nieri-Bastos<sup>1</sup>, Márcia Aparecida Sperança<sup>3,4</sup> and Arlei Marcili<sup>1,5\*</sup>

## Abstract

**Background:** *Leishmania infantum*, the etiological agent of visceral leishmaniasis, is a neglected zoonosis that requires validation and standardization of satisfactory diagnostic methodologies. Thus, the aim of the present study was to evaluate the effectiveness of cathepsin L-like protease as a target for making molecular diagnoses and as a phylogenetic marker enabling to understand the intraspecies variations and evolutionary history of *L. infantum* in Brazil.

**Methods:** We used 44 isolates of *L. infantum*. The cathepsin L-like gene fragments were amplified, sequenced, manually aligned and analyzed using inference methods. The sequences generated were used to search and design oligonucleotide primers to be used in reactions specific to the target parasite.

**Results:** The cathepsin L-like gene did not show any intraspecies variability among the isolates analyzed. The pair of primers proposed amplified the target deoxyribonucleic acid (DNA) of *L. infantum* isolates and were effective for DNA amplification at concentrations of as low as  $10^{-11}$  ng/ $\mu$ l. The proposed marker did not present cross-reactions with other hemoparasites. When used for making the diagnosis in a panel of clinical samples from dogs, a positivity rate of 49.03% (102/208) was obtained, versus 14.42% (30/208) for a ribosomal internal transcribed spacer (ITS) marker. In samples from sandflies, the rate was 6.25% and from humans, 14.28%.

**Conclusions:** The results described in this work allow us to infer that CatLeish-PCR is a sensitive and specific marker for use in diagnostic trials of *L. infantum* and in clinical and epidemiological surveys.

**Keywords:** *Leishmania (Leishmania) infantum*, Cysteine proteases, Cathepsin L-like, Molecular diagnosis

## Background

*Leishmania (Leishmania) infantum* is a flagellate protozoan with a heterogenic cycle that belongs to the genus *Leishmania*, which is in the family Trypanosomatidae of the order Kinetoplastida [1]. This parasite is the etiological agent for visceral leishmaniasis, a zoonosis with worldwide distribution. Most of the cases are

concentrated in Bangladesh, Nepal, India, South Sudan and Brazil. The development of this parasite comprises two morphological forms, whose main differences are the position of the kinetoplast in relation to the nucleus and the presence of a free flagellum associated with the undulating membrane [2, 3]. The promastigote form is found in arthropod vectors, represented by insects of the phlebotomine group [4], while amastigote forms are found inside cells of the mononuclear phagocytic system of vertebrate hosts such as rodents, marsupials, bats, canids and humans [5, 6].

The severity of the disease is explained by the high virulence of its etiological agents, which results from the

\* Correspondence: [amarcili@usp.br](mailto:amarcili@usp.br)

<sup>1</sup>Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, Av. Prof. Dr. Orlando Marques de Paiva, 87, São Paulo, SP 05508-270, Brazil

<sup>5</sup>Medicina Veterinária e Bem estar animal, Universidade Santo Amaro, São Paulo, SP, Brazil

Full list of author information is available at the end of the article



action of a set of functional molecules with intense biological activity in their hosts. The role of protease inhibitors in the development of pathogenic mechanisms was investigated [7]. These enzymes can be divided into endopeptidases and exopeptidases, depending on the site of hydrolysis. If the residue present at the catalytic site is taken into account, they are divided into metalloproteases, serine proteases, aspartic proteases and cysteine proteases [8]. In the genus *Leishmania*, cysteine proteases are the most abundant class of enzymes and are concentrated inside megasomes, where they act towards regulation of metabolic routes, cell differentiation in vertebrates and vectors, cell invasion and transposition of tissue barriers, degradation of hemoglobin and other hematopoietic proteins, evasion of the immune response, activation of inflammatory responses and programmed cell death [9].

Cathepsin L-like cysteine proteases in *Leishmania* have three isoforms, named cysteine protease A (CPA), cysteine protease B (CPB) and cysteine protease C (CPC). These are biochemically organized into four domains (predomain, prodomain, catalytic domain, C-terminal extension) that result from expression of a multigenic family arranged in tandem. In the specific case of *L. infantum*, two isoforms are expressed, the CPB isoform expressed in the promastigote forms found in the insect vectors and the CPA isoform, in which transcriptomic studies revealed a unique expression profile of the amastigote forms [10].

They are, therefore, used to construct phylogenetic inferences of close sequences and to resolve problems regarding polytomy and inferences of low support [11, 12]. Cathepsin genes have already been used for understanding phylogenetic relationships and as a target for making molecular diagnoses regarding other trypanosomatid species [13–15]. However, there are no studies that characterize this gene in *L. infantum* or that investigate it as a possible diagnostic marker that might help to solve recurrent problems regarding diagnostic investigation of this parasite within the clinical routine and in epidemiological investigations.

Despite the high importance of this disease, there is some difficulty in standardizing diagnostic methodologies with high predictive values for reservoir surveys. Making a direct diagnosis is invasive and laborious, and only low levels of sensitivity are reached [16–18]. The serological tests also have a series of technical limitations, such as low specificity values resulting from cross-reactions with other trypanosomatids, low concordance indices between different serological tests and lack of consensus regarding the nature and use of the antigenic product to be employed [19–22].

Thus, the objective of this study was to evaluate the CPA isoform of cathepsin L-like sequences as a marker for genetic analysis on intraspecific variability of *L.*

*infantum* and as a marker for making molecular diagnoses on visceral leishmaniasis.

## Methods

### Leishmania isolates, DNA preparation, amplification and sequencing of cathepsin L-like gene

DNA from 44 *Leishmania* isolates (Table 1) was extracted from culture supernatants using the phenol-chloroform method and from primary samples (human blood, urine, conjunctival swabs from dogs and sandfly material) in accordance with the protocol established for the Purelink kit (Thermo Fisher Scientific Inc., 2012, USA).

First of all, the extracted DNA samples were quantified and submitted to conventional polymerase chain reactions (PCR) using the specific primers to Mammalian Cytochrome B as a constitutive gene, to ensure the quality of the samples [23]. After this quality control, the DNA samples were submitted to the PCR reactions using high-fidelity Taq DNA polymerase with the specific primers designated for Cathepsin L-like CPA from *Leishmania* [24] which comprised a fragment of around 893 base pairs (bp).

All the isolates were included in the Brazilian Trypanosomatid Collection (Coleção Brasileira de Tripanossomatídeos, CBT) of the School of Veterinary Medicine of the University of São Paulo, Brazil.

### Phylogenetic analysis

The sequences obtained were aligned with sequences retrieved from GenBank using ClustalX [25] and were adjusted manually using GeneDoc [26] and then deposited in GenBank (Table 1). The cathepsin L-like CPA sequences were used to construct a phylogenetic tree using maximum parsimony, as implemented in PAUP version 4.0b10 [27] with 500 bootstrap replicates. Bayesian analysis was performed using MrBayes v3.1.2 [28] with 1,000,000 replicates. The first 25% of the trees represented burn-in, and the remaining trees were used to calculate Bayesian posterior probability.

### Standardization of *L. infantum*-specific assay based on cathepsin L-like protease

The aligned cathepsin L-like CPA gene sequences were used to search for consensus regions and to design specific primers for diagnosing *L. infantum*. The criteria used to define the primer pair were the guanine-to-cytosine ratio, formation of guanidine/cytosine (GC) staples, formation of self-homologies, self-dimer formation measured using the  $\Delta G$  value, melting temperature and in silico specificity of the primers through BLASTn.

A specific PCR procedure (designated CatLeish-PCR) was developed for amplification of 223 bp of genomic DNA from *L. infantum*, using the CatLeishF primer (5' GACAACGGCACCGTCGGCGCCAAAAT

**Table 1** *Leishmania infantum* isolates, host, geographical origin and sequences of Cathepsin L-like employed in the phylogenetic analysis performed in this study

CBT <sup>a</sup>	Host	Geographical origin	Accession number <sup>b</sup>
01	<i>Canis familiaris</i>	SP	MH427793
12	<i>Canis familiaris</i>	Recreio	MH427794
13	<i>Canis familiaris</i>	Mangaratiba	MH427795
14	<i>Canis familiaris</i>	Ilha Grande	MH427796
15	<i>Canis familiaris</i>	Barra da Tijuca	MH427797
16	<i>Canis familiaris</i>	Recreio	MH427798
17	<i>Canis familiaris</i>	Cuiabá	MT
18	<i>Canis familiaris</i>	Ilha de Guaratiba	RJ
20	<i>Canis familiaris</i>	Cuiabá	MT
22	<i>Canis familiaris</i>	Caucaia	CE
23	<i>Canis familiaris</i>	Fortaleza	CE
24	<i>Canis familiaris</i>	Jequié	BA
25	<i>Canis familiaris</i>	Campo Grande	MS
26	<i>Canis familiaris</i>		DF
27	<i>Cerdocyon thous</i>		PA
28	<i>Canis familiaris</i>	Teresina	PI
29	<i>Canis familiaris</i>	Teresina	PI
30	<i>Canis familiaris</i>	Uruguaiana	RS
31	<i>Canis familiaris</i>	Uruguaiana	RS
34	<i>Canis familiaris</i>	Petrolina	PE
37	<i>Canis familiaris</i>	Petrolina	PE
39	<i>Canis familiaris</i>	Santarém	PA
40	<i>Canis familiaris</i>	Santarém	PA
43	<i>Canis familiaris</i>	Santarém	PA
44	<i>Canis familiaris</i>	Santarém	PA
49	<i>Canis familiaris</i>	Campo Grande	MS
50	<i>Canis familiaris</i>	Campo Grande	MS
54	<i>Canis familiaris</i>	Teresina	PI
55	<i>Canis familiaris</i>	Teresina	PI
56	<i>Canis familiaris</i>	Campo Grande	MS
57	<i>Canis familiaris</i>	Campo Grande	MS
62	<i>Canis familiaris</i>	Petrolina	PE
96	<i>Canis familiaris</i>	Caxias	MA
105	<i>Canis familiaris</i>	Natal	RN
106	<i>Canis familiaris</i>	Natal	RN
107	<i>Canis familiaris</i>	Natal	RN
108	<i>Canis familiaris</i>	Natal	RN
124	<i>Canis familiaris</i>	Natal	RN
125	<i>Canis familiaris</i>	Natal	RN
126	<i>Canis familiaris</i>	Natal	RN
153	<i>Canis familiaris</i>	São Domingos	MA
179	<i>Canis familiaris</i>	Patos	PB

**Table 1** *Leishmania infantum* isolates, host, geographical origin and sequences of Cathepsin L-like employed in the phylogenetic analysis performed in this study (Continued)

CBT <sup>a</sup>	Host	Geographical origin	Accession number <sup>b</sup>
211	<i>Homo sapiens</i>	Marília	SP
212	<i>Homo sapiens</i>	Marília	SP

Brazilian states: BA, Bahia; CE, Ceará; DF, Distrito Federal; MA, Maranhão; MS, Mato Grosso do Sul; MT, Mato Grosso; PA, Pará; PB, Paraíba; PE, Pernambuco; PI, Piauí; RJ, Rio de Janeiro; RN, Rio Grande do Norte; RS, Rio Grande do Sul; SP, São Paulo

<sup>a</sup>Coleção Brasileira de Tripanossomatídeos

<sup>b</sup>GenBank accession number

AAAAG 3') and CatLeishR primer (5' CAGTAC GGCGGTTTCGCTTGTTGAAGC 3') (Fig. 1). The standard conditions for amplification of the cathepsin L-like CPA sequences comprised 34 cycles of denaturation at 94 °C for 1 min, annealing at 64 °C for 1 min and extension at 72 °C for 45 s.


#### Sensitivity and specificity assay on CatLeish-PCR

For sensitivity tests, DNA from *L. infantum* was serially diluted at the concentrations of  $1 \times 10^{-7}$ ,  $1 \times 10^{-8}$ ,  $1 \times 10^{-9}$ ,  $1 \times 10^{-10}$ ,  $1 \times 10^{-11}$ ,  $1 \times 10^{-12}$ ,  $1 \times 10^{-13}$ ,  $1 \times 10^{-14}$  and  $1 \times 10^{-15}$  ng/μl.

Specificity tests were performed on DNA samples from other parasite species in the genus *Leishmania*, including: *Leishmania (Viannia) braziliensis*, *Leishmania (Viannia) guyanensis*, *Leishmania (Viannia) naiffi*, *Leishmania (Leishmania) amazonensis*, *Leishmania (Leishmania) mexicana*, *Leishmania (Sauroleishmania) gymnodactyli*, *Leishmania (Sauroleishmania) adleri*, *Leishmania (Sauroleishmania) tarentolae* and *Leishmania (Mundinia) enriettii*. In addition, the following species in the genus *Trypanosoma* were tested: *Trypanosoma dionisi*, *Trypanosoma terrestris*, *Trypanosoma cruzi*, *Trypanosoma cruzi marinkellei*, *Trypanosoma theileri* and *Trypanosoma gennarii*; along with two other pathogenic species that are common in dogs: *Babesia canis* and *Ehrlichia canis*.

#### Application of CatLeish-PCR to clinical samples

CatLeish-PCR was tested on a panel of clinical samples of purified DNA obtained from dogs in endemic and non-endemic regions in São Paulo state, Brazil. In addition, blood samples (prepared on filter paper), conjunctival swabs and urine samples from dogs that were known to be positive in the parasitological test were used. DNA samples obtained from human patients blood from Marília, São Paulo state and DNA extracted from sandflies caught in Bom Jesus dos Perdões, São Paulo state were also tested in pools. All samples were further tested using the ITS rDNA marker [29].



**Fig. 1** Alignment of cathepsin L-like CPA sequences from *Leishmania infantum*. Schematic representation of primers used for PCR amplification

## Results

Forty-four sequences were obtained from cathepsin L-like CPA and they were all identical, without polymorphism, and presented 99% similarity with cysteine peptidase isoform A of *Leishmania infantum* (XM\_001465076.1) from Europe. The sequences of cathepsin L-like CPA from *L. infantum* were identical and all isolates were clustered together (100% bootstrap/100% posterior probability and 100% similarity) (Fig. 2).

*Leishmania* species were aggregated in a monophyletic group (Fig. 2). The different species of the genus *Leishmania* were segregated into four groups: I. *Leishmania* species with mucocutaneous clinical manifestations, including *L. guyanensis*, *L. braziliensis* and *L. panamensis* (0.1% divergence of sequences and 100% bootstrap and 1.0 posterior probability); II. Species with cutaneous manifestations, including *L. pifanoi*, *L. mexicana* and *L. amazonensis* (0.52% divergence of sequences and 100% bootstrap and 1.0 posterior probability); III. *Leishmania* species causing cutaneous “oriental sore”, including *L. major* and *L. aethiopica* (1.53% divergence of sequences and 100% bootstrap and 1.0 posterior probability); and IV. New and Old-World species of visceral leishmaniasis (0.14% divergence of sequences and 100% bootstrap and 1.0 posterior probability), separated according to the geographical origin of the isolates (Fig. 2).

The in silico analyses on the primers proposed for the specific PCR did not indicate any possibility of formation of GC staples, self-dimers and auto-homologies. The CatLeishF primer showed a GC ratio of 54.5% with an estimated melting temperature of 66.2 °C, whereas the CatLeishR primer had a GC ratio of 57.6% with a melting temperature corresponding to 65.5 °C. In BLASTn analysis, these primers were specific for *L. infantum*,


with no homology containing sequences from other species deposited in the database.

CatLeish-PCR reactions with genomic DNA from *L. infantum* isolates demonstrated that there was a correct amplification, since amplicon sequencing revealed in BLASTn a percentage of 99% similarity to deposited sequences of cysteine protease genes from European strains of *L. infantum* (XM\_001465076.1) and identical to American *L. infantum* sequences obtained in this study. Sensitivity measurement demonstrated that the markers were efficient at amplifying the target DNA and forming detectable bands at concentrations of up to  $10^{-11}$  ng/μl (Fig. 3).

Different species of the genus *Leishmania* were tested and there was no cross-amplification with any of the species epidemiologically associated with the cutaneous or tegumentary leishmaniasis complex in Brazil. When the molecular diagnosis was tested against hemoparasites belonging to the genera *Babesia*, *Ehrlichia* and *Trypanosoma*, there was no amplification (Fig. 3).

The use of CatLeish-PCR in the panel of clinical samples revealed that frequency of positive dogs was 49.03%, thus contrasting with the 14.42% achieved through using primers for the ribosomal ITS gene. All samples that were positive in the reactions for the ITS gene were positive in reactions for the cathepsin gene. In the human samples, 6.25% were positive, whereas in samples from sandflies, the positivity was 14.28% in pools (Table 2).

The samples from parasitologically positive dogs showed different rates of positivity according to the material tested. Urine samples showed 33% positivity, conjunctival swabs 50%, samples preserved on filter paper 100% and whole blood samples 100%.




The CatLeish-PCR was replicated several times during standardization with DNA samples from American *L. infantum* isolates, and subsequently technical replication was tested with clinical samples. Regardless of the samples tested, the technical replication and amplification standard were maintained (Additional file 1).

## Discussion

The phylogenetic inferences from maximum parsimony and Bayesian analyses revealed a pattern of high similarity for *L. infantum* isolates, as observed in

inferences based on SSUrDNA and gGAPDH genes [30]. The segregation of *L. infantum* from Europe and America corroborates the notion that this species was introduced to the Americas during the colonization period [30].

This absence of divergence, even among isolates from biomes with different climates, rainfall, vegetation cover and availability of hosts and arthropod vectors, corroborates the theory that this parasite is not autochthonous to the Americas [31]. This finding indicates that the event leading to introduction of this parasite to the New



**Fig. 3** CatLeish-PCR for diagnosis of *Leishmania infantum* based on Cathepsin L-like sequences. **a** Specificity analysis using DNA, *Leishmania* and *Trypanosoma* sequences and other hemoparasites from dogs (*Babesia* and *Ehrlichia*). **b** Sensitivity analysis using DNA from *Leishmania infantum* in different concentrations

World was recent. It can therefore probably be dated to the process of Ibero-American colonization that began in the fifteenth century, caused by the arrival of infected dogs and rodents [32]. Focusing only on a few specific genes we can speculate that this short evolutionary time interval would be insufficient for the appearance and fixation of possible mutations in the population.

These results refute the hypothesis of a scenario in which the parasite responsible for cases of visceral leishmaniasis is native to the Americas [30–33]. Thus, the argument that the parasite is autochthonous in that it is adapted to parasitism in other wild fauna reservoirs can be countered through attributing this adaptation to the eclectic dietary habits of the vector insects [34].

The phylogenetic analysis on the cathepsin L-Like CPA gene showed clustering that reflects the distinct

clinical manifestations of *Leishmania* infection, and it was similar to amino acid analyses on cysteine proteases [24]. In addition, the topology generated using the cathepsin L-like gene also corroborates data generated using ribosomal spacers and cytochrome B [35, 36].

The inferences correctly solved the differentiation of the species that are considered to be the etiological agent of tegumentary forms of the disease. One of the clades grouped *L. guyanensis*, *L. panamensis* and *L. braziliensis*, and this reflected the findings of studies in which, analyzing otorhinolaryngological conditions, most of the etiology of mucocutaneous leishmaniasis was attributed to *L. braziliensis* and *L. panamensis*, as well as results in which clinical situations of mucosal involvement were attributed to *L. guyanensis* [37, 38].

A second group included *L. amazonensis* and *L. mexicana*, species that have been widely associated with cutaneous manifestations of tegumentary leishmaniasis in the New World [39–43]. The *L. pifanoi* sequences were positioned in this same group. Although this species has lower epidemiological impact regarding disease transmission, it belongs to the *L. mexicana* complex [44]. Therefore, this result corroborates the data demonstrating similarity among the species of this complex, with regard to proposing specific markers for kinetoplast DNA (kDNA) [45].

These results confirm that the cathepsin L-like CPA gene is a good marker for phylogenetic positioning of species of the genus *Leishmania*, which suggests that this gene may be a good target for making the molecular diagnosis of

**Table 2** Positivity of naturally infected biological samples based on Cathepsin L-like of *Leishmania infantum* (CatLeish-PCR) marker

Samples (number)	Positives samples (%)	
	CatLeish-PCR	ITS
Dogs blood (208)	49,03 (102)	14,42 (30)
Dogs urine (3)	33 (1)	
Dogs conjunctival swab (4)	50 (2)	
Dogs blood in filter paper (4)	100 (4)	
Human blood (50)	6,25 (3)	
Sand flies (21)	14,28 (3)	

leishmaniasis. This would enable satisfactory differentiation between the clinical forms of the disease, including among the variants of the tegumentary forms, which may aid in prescribing therapy, establishing the medical prognosis for the disease and mitigating the recurrent problems of specific diagnosis. We agree that the number of isolates collected in this work is sufficient to portray the dispersal of the parasite in Brazil.

Cathepsin gene-based assays have shown excellent results with regard to diagnosing the following *Trypanosoma* species: *T. vivax* [13], *T. congolense* [46], *T. theileri* [47], *T. cruzi* [48] and *T. rangeli* [49]. However, the low numbers of copies of this gene in the genome are very specific and especially sensitive [13].

Cathepsin L-like protease is differentially expressed in different forms of the parasite [50, 51]. The CPA isoform is preferentially expressed in amastigote forms. In vertebrates (both human and animal hosts), amastigotes are the replicative form and cathepsin L-like CPA is continuously expressed [24]. The expression pattern of cathepsin L-like CPA and the presence of immunogenic epitopes may indicate it as an interesting target in diagnostic-serological tests, which are preferentially used in the clinical routine [25].

The diagnostic marker based on the cathepsin L-like CPA gene has shown high sensitivity and was specific for *L. infantum*, which enables direct use in clinical samples both in non-endemic areas with imported cases and in endemic areas in which different species of *Leishmania* may be circulating. Thus, it has been proven to be effective for epidemiological surveys on human hosts, animal reservoirs and arthropod vectors. Another advantage that makes CatLeish-PCR feasible is that there is no need for complementary restriction enzyme digestion steps or use of robust high-resolution melting equipment for identifying *L. infantum*, which makes the method less costly and less cumbersome than other methodologies that are available [52].

In comparison with the markers for the ITS gene, CatLeish-PCR demonstrated higher prevalence of leishmaniasis, which thus corroborated the diagnostic sensitivity of the marker. The marker was effective in making the diagnosis, both from positive swab samples from conjunctival lesions and from blood samples fixed on filter paper, reinforcing the high sensitivity of CatLeish-PCR. The positive animals in the parasitological examination were also positive when submitted to molecular diagnosis based on the cathepsin L-like gene, except for the urine samples given the low amount of DNA in two animals. This versatility is important because samples fixed on filter paper are easier to transport and store, given their stability, as well as exempt invasive collection methodologies [29].

The diagnostic method works very well for blood samples and even though the number of samples from other sources is reduced, the method borrows our hypothesis

of good molecular marker for diagnosis in biological samples with DNA. Amplification was observed even in samples extracted from urine, whose DNA concentration is considerably reduced, corresponding only to fragments of the dead parasite excreted in the urinary system [53]. It should be noted that even with low levels of positivity in the assays using biological samples as a template, we still support our indication of the use of the Cathepsin gene for PCR diagnosis, since even in biological samples where the genetic material of the parasites is scarce or damaged, we obtained satisfactory amplifications. Considering this, CatLeish-PCR may have potential for monitoring the remission of infection in patients with visceral leishmaniasis and as an indicator of the efficacy of possible therapeutic approaches [54].

The results presented here allow us to propose that CatLeish-PCR is a tool that can be used for diagnosing visceral leishmaniasis. It surmounts the recurrent problems of low sensitivity that direct visualization methodologies present and those of low specificity and agreement among the serological methods that have preferentially been recommended for diagnosing this disease [55]. The molecular diagnostic assays used and described in the literature have low sensitivity and specificity. Markers based on the cytochrome b gene are non-specific [56]. Markers based on genes present in the kinetoplast have high sensitivity, but their specificity is low [57, 58]. Markers based on ribosomal genes are specific, but not very sensitive [59]. Other assays such as restriction fragment length polymorphism (PCR-RFLP) have been standardized, but these raise the cost of making the diagnosis or still require sequencing of the product [60–62].

Epidemiological studies are important for recognizing active transmission cycles or introduction of parasites into new areas. Therefore, precise diagnostic methods are essential for diagnosing the presence of a parasite that is still emerging in many regions. Moreover, making a precise diagnosis minimizes the need to destroy dogs, which are the main reservoirs, given that in some countries this prophylactic measure used in combating human leishmaniasis.

## Conclusion

This is the first work that characterizes the CPA isoform of the L-like cathepsin gene of *L. infantum*, demonstrating the lack of genetic variability among the Brazilian isolates of *L. infantum*. We propose that the gene studied here be an efficient phylogenetic marker for parasites of the genus *Leishmania*, and also the developed CatLeish-PCR has been shown to be sensitive and specific for the effective clinical diagnosis of this zoonosis in dogs and humans.

## Supplementary information

**Supplementary information** accompanies this paper at <https://doi.org/10.1186/s12879-019-4463-8>.

**Additional file 1.** Amplification patterns of CatLeish-PCR for diagnosis of *Leishmania infantum* in samples from dogs, humans and sand flies.

### Abbreviations

C: Cytosine; CBT: Coleção Brasileira de Tripanossomatídeos; CPA: Cysteine protease A; CPB: Cysteine protease B; CPC: Cysteine protease C; DNA: Deoxyribonucleic acid; G: Guanidine; ITS: Internal transcribed spacer; kDNA: kinetoplast deoxyribonucleic acid; PCR: Polymerase chain reaction; PCR-RFLP: Restriction fragment length polymorphism

### Acknowledgments

We would like to thank to veterinary Vilma Pereira da Costa and all the people who assisted this study in the different locations studied.

### Authors' contributions

RES and AM conceived the study and designed the experiments; RES, BMS, LESC, APC, FNB and MAS, assisted with sample collection, performed the experiments and analyzed the data; RES, RT and AM prepared the paper. All authors read and approved the final version of the manuscript.

### Funding

This research was financially supported by Fundação de Amparo a Pesquisa do Estado de São Paulo (FAPESP) process number 2010/50886-7, and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) process number 404249/2016-7 and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES). RES scholarship from CNPq, BMC from CAPES and APC from FAPESP,

### Availability of data and materials

The accession numbers of GenBank from sequences obtained in this study are described in Table 1.

### Ethics approval and consent to participate

The human samples were transferred by the Laboratory of Pathogens from Center of Natural and Human Sciences from Universidade Federal do ABC and approved by Ethics Committee for Human research from Federal University of ABC (number 883722718900005594). The clinical samples from dogs and sandflies procedures were endorsed by the Ethics Committee for Animal Use of the University of São Paulo, Brazil (number 7584250216) according National Council for Control of Animal Experimentation.

### Consent for publication

Not applicable.

### Competing interests

The authors declare that they have no competing interests.

### Author details

<sup>1</sup>Departamento de Medicina Veterinária Preventiva e Saúde Animal, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, Av. Prof. Dr. Orlando Marques de Paiva, 87, São Paulo, SP 05508-270, Brazil.

<sup>2</sup>Faculdade de Medicina, Universidade Santo Amaro, São Paulo, SP, Brazil.

<sup>3</sup>Ciência Animal, Universidade Estadual do Maranhão, São Luís, MA, Brazil.

<sup>4</sup>Centro de Ciências Naturais e Humanas, Universidade Federal do ABC, São Bernardo do Campo, SP, Brazil. <sup>5</sup>Medicina Veterinária e Bem estar animal, Universidade Santo Amaro, São Paulo, SP, Brazil.

Received: 21 February 2019 Accepted: 11 September 2019

Published online: 28 October 2019

### References

- Honigberg BM. A contribution to systematics of the non-pigmented flagellates. In: Ludvík J, Lom J, Vavra J, editors. Progress in Protozoology. New York: Academic Press; 1963. p. 68–9.
- Wallace FG. Biology of the Kinetoplastida of Arthropods. In: Lumsden WHR, Evans DA, editors. Biology of the Kinetoplastida. New York: Academic Press; 1979. p. 213–40.
- Vickerman K. The evolutionary expansion of the trypanosomatid flagellates. Int J Parasitol. 1994;24:1317–31.
- Galvis-Ovallos F, Silva MD, Bispo GB, Oliveira AG, Neto JR, Malafronte RD, et al. Canine visceral leishmaniasis in the metropolitan area of São Paulo: *Pintomyia Fischeri* as potential vector of *Leishmania infantum*. Parasite. 2017;24:2.
- Deane LM, Deane MP. Encontro de *Leishmania* nas vísceras e na pele de uma raposa, em zona endêmica de calazar, nos arredores de Sobral, Ceará. Hospital. 1954;45:419–21.
- Costa AP, Costa FB, Soares HS, Ramirez DG, Mesquita ET, Gennari SM, et al. *Trypanosoma cruzi* and *Leishmania infantum chagasi* infection in wild mammals from Maranhão state, Brazil. Vector Borne Zoonotic Dis. 2015;15:656–66.
- Gomes CB, Silva FS, Charret KS, Pereira BAS, Finkelstein LC, Souza RS, et al. Increasing in cysteine proteinase B expression and enzymatic activity during in vitro differentiation of *Leishmania (Viannia) braziliensis*: first evidence of modulation during morphological transition. Biochimie. 2016;133:28–36.
- Silva-López RE. Proteases de *Leishmania*: novos alvos para o desenvolvimento racional de fármacos. Quím Nova. 2010;33:1541–8.
- Agallou M, Margaroni M, Athanasiou E, Toubanaki D, Kontonikola K, Karidi K, et al. Identification of BALB/c immune markers correlated with a partial protection to *Leishmania infantum* after vaccination with a rationally designed multi-epitope cysteine protease a peptide-based nanovaccine. PLoS Negl Trop Dis. 2017;11:1.
- Mundodi V, Kucknoor AS, Gedamu L. Role of *Leishmania (Leishmania) chagasi* amastigote cysteine protease in intracellular parasite survival: studies by gene disruption and antisense mRNA inhibition. BMC Mol Biol. 2005;6:3.
- Mottram JC, Coombs GH, Alexander J. Cysteine peptidases as virulence factors of *Leishmania*. Curr Opin Microbiol. 2004;7:375–81.
- Casgrain PA, Martel C, McMaster WR, Mottram JC, Olivier M, Descoteaux A. Cysteine peptidase B regulates *Leishmania mexicana* virulence through the modulation of GP63 expression. PLoS Pathog. 2016;12:5.
- Cortez AP, Rodrigues AC, Garcia HA, Neves L, Batista JS, Bengaly Z, et al. Cathepsin L-like genes of *Trypanosoma vivax* from Africa and South America characterization, relationships and diagnostic implications. Mol Cell Probes. 2008;23:44–51.
- Garcia HA, Kamyingkird K, Rodrigues AC, Jittapalapong S, Teixeira MM, Desquesnes M. High genetic diversity in field isolates of *Trypanosoma theileri* assessed by analysis of cathepsin L-like sequences disclosed multiple and new genotypes infecting cattle in Thailand. Vet Parasitol. 2011;180:363–7.
- Yokoyama N, Sivakumar T, Fukushi S, Tattiyapong M, Tuvshintulga B, Kothalawala H, et al. Genetic diversity in *Trypanosoma theileri* from Sri Lankan cattle and water buffaloes. Vet Parasitol. 2015;207:335–41.
- Ferrer L. Clinical aspects of canine leishmaniasis. In: R Killick-Kendrick, editor. Canine leishmaniasis: an update. Proceedings of a Canine Leishmaniasis Forum. Barcelona: Hoeschst Roussel Vet; 1999. p. 6–10.
- Assis TSM, Braga ASC, Pedras MJ, Barral AMP, Siqueira IC, Costa CHN, et al. Validação do teste imunocromatográfico rápido IT-LEISH® para o diagnóstico da leishmaniose visceral humana. Epidemiol Serv Saúde Brasília. 2008;17:107–16.
- Laurenti MD. Correlação entre o diagnóstico parasitológico e sorológico na leishmaniose visceral americana canina. Bol Epidemiol Paul. 2009;6:13–23.
- Burns JM, Shreffler WG, Benson DR, Ghalib HW, Badaro R, Reed SG. Molecular characterization of a kinesin-related antigen of *Leishmania chagasi* that detects specific antibody in African and American visceral leishmaniasis. Proc Natl Acad Sci. 1993;90:775–9.
- Mancianti F, Pedoneze F, Poli A. Evaluation of dot enzyme-linked immunosorbent assay (dot-ELISA) for the serodiagnosis of canine leishmaniasis as compared with indirect immunofluorescence assay. Vet Parasitol. 1996;65:1–9.
- Ferroglio E, Centaro E, Mignone W, Trisciuglio A. Evaluation of an ELISA rapid device for the serological diagnosis of *Leishmania infantum* infection in dog as compared with immunofluorescence assay and Western blot. Vet Parasitol. 2006;144:162–6.
- Hirschmann LC, Brod CS, Radin J, Simon CF, Recuero AL. Leishmaniose visceral canina: comparação de métodos sorológicos em cães de área indígena do Rio Grande do Sul no Brasil. Rev Patol Trop. 2015;44:33–44.

23. Yanh BB, Chen DL, Chen JP, Liao L, Hu XS, Xu JN. Analysis of kinetoplast cytochrome b gene of 16 Leishmania isolates from different foci of China: different species of *Leishmania* in China and their phylogenetic inference. *Parasit Vectors.* 2013;6:32.
24. Kuru T, Jirata D, Genetu A, Barl S, Mengistu Y, Aseffa A, et al. *Leishmania aethiopica*: identification and characterization of cathepsin L-like cysteine protease genes. *Exp Parasitol.* 2007;115:283–90.
25. Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The clustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 1997;25:4876–82.
26. Nicholas KB, Nicholas HB, Deerfield DW, Nicholas HBJ, DWID, Nicholas KB, et al. GeneDoc: analysis and visualization of genetic variation. *Emblnewsm.* 1997;4:24.
27. Swofford DL. PAUP: phylogenetic analysis using parsimony. Version 4b6. Sunderland: Sinauer Associates; 1998.
28. Huelsenbeck JP, Ronquist F. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics.* 2001;17:754–5.
29. El Tai NO, Osman OF, El Fari M, Presber W, Schönian G. Genetic heterogeneity of ribosomal internal transcribed spacer in clinical samples of *Leishmania donovani* spotted on filter paper as revealed by single-strand conformation polymorphisms and sequencing. *Trans R Soc Trop Med Hyg.* 2000;94:575–9.
30. Marcili A, Sperança MA, Costa AP, Madeira MF, Soares HS, Sanches CO, et al. Phylogenetic relationships of *Leishmania* species based on trypanosomatid barcode (SSU rDNA) and gGAPDH genes: taxonomic revision of *Leishmania (L.) infantum chagasi* in South America. *Infect Genet Evol.* 2014;25:44–51.
31. Akhouni M, Kuhls K, Cnnett A, Votypka J, Marty P, Delaunay P, et al. A historical overview of the classification, evolution, and dispersion of *Leishmania* parasites and sandflies. *PLoS Negl Trop Dis.* 2016;10:3.
32. Mauricio IL, Stothard JR, Miles MA. The strange case of *Leishmania chagasi*. *Parasitol Today.* 2000;16:188–9.
33. Lainson R, Shaw JJ. Evolution, classification and geographic distribution. In: Peters W, Killick-Kendrick R, editors. *The Leishmaniasis in biology and medicine*. London: Academic Press; 1978. p. 1–120.
34. Travi BL, Osorio Y, Becerra MT, Adler GH. Dynamics of *Leishmania chagasi* infection in small mammals of the undisturbed and degraded tropical dry forests of northern Colombia. *Trans R Soc Trop Med Hyg.* 1998;92:275–8.
35. Dávila AM, Momen H. Internal-transcribed-spacer (ITS) sequences used to explore phylogenetic relationships within *Leishmania*. *Ann Trop Med Parasitol.* 2000;94:651–4.
36. Luyo-Acero GE, Uezato H, Oshiro M, Takei K, Kariya K, Katakura K, et al. Sequence variation of the cytochrome b gene of various human infecting members of the genus *Leishmania* and their phylogeny. *Parasitology.* 2004;128:483–91.
37. Fornazieri MA, Yamaguti HY, Moreira JH, Takemoto LE, Navarro PL, Hesbke RE. The Most common Otorhinolaryngologic manifestations of granulomatous diseases. *Int Arch Otorhinolaryngol.* 2008;12:362–5.
38. Souza IS, Naiff RD, Guimarães TC, Naiff MF, Cupolilo E, Rodrigues WA, et al. American cutaneous leishmaniasis due to *Leishmania (Viannia) guyanensis* as an initial clinical presentation of human immunodeficiency virus infection. *J Eur Acad Dermatol Venereol.* 1998;10:214–7.
39. Seidelin H. Leishmaniasis and Babesiosis in Yucatan. *Ann Trop Med Parasitol.* 1912;6:295–300.
40. Silveira FT, Lainson R, Shaw JJ, Souza AA, Ishikawa EA, Braga RR. Cutaneous leishmaniasis due to *Leishmania (Leishmania) amazonensis* in Amazonian Brazil, and the significance of a negative Montenegro skin-test in human infections. *Trans R Soc Trop Med Hyg.* 1991;85:735–8.
41. Martinez E, Le Pont F, Torrez M, Telleria J, Vargas F, Muñoz M, et al. A new focus of cutaneous leishmaniasis due to *Leishmania amazonensis* in a sub Andean region of Bolivia. *Acta Trop.* 1998;71:97–106.
42. Narvaez FJA, Medina-Peralta S, Vargas-Gonzalez A, Canto-Lara SB, Estrada-Parra S. The histopathology of cutaneous leishmaniasis due to *Leishmania (Leishmania) Mexicana* in the Yucatan peninsula, Mexico. *Rev Inst Med Trop Sao Paulo.* 2005;47:191–4.
43. Lindoso JAL, Cunha MA, Queiroz IT, Moreira CHV. Leishmaniasis–HIV coinfection: current challenges. *HIV AIDS (Auckl).* 2016;8:147–56.
44. Colmenares M, Constant SL, Kima PE, McMahon-Pratt D. *Leishmania pifanoi* pathogenesis: selective lack of a local cutaneous response in the absence of circulating antibody. *Infect Immun.* 2002;70:6597–605.
45. Montes OH, Guzmán SG, Gómez FM, Barker DC, Monroy-Ostria A. Analysis of Kinetoplast DNA from Mexican isolates of *Leishmania (L.) mexicana*. *Interdiscip Perspect Infect Dis.* 2012. <https://doi.org/10.1155/2012/279081>.
46. Mendonça-Palomares C, Biteau N, Giroud C, Coustou V, Coetzter T, Authié E, et al. Molecular and biochemical characterization of a Cathepsin B-like protease family unique to *Trypanosoma congolense*. *Eukaryot Cell.* 2008;7:684–97.
47. Rodrigues AC, Garcia HA, Ortiz PA, Cortez AP, Martinkovic F, Paiva F, et al. Cysteine proteases of *Trypanosoma (Megatrypanum) theileri*: cathepsin L-like gene sequences as targets for phylogenetic analysis, genotyping diagnosis. *Parasitol Int.* 2010;58:318–25.
48. Lima L, Ortiz PA, Silva FM, Alves JM, Serrano MG, Cortez AP, et al. Repertoire, genealogy and genomic organization of cruzipain and homologous genes in *Trypanosoma cruzi*, *T. cruzi-like* and other trypanosome species. *PLoS One.* 2012;7:6.
49. Ortiz PA, Maia da Silva F, Cortez AP, Lima L, Campaner M, Pral EM, et al. Genes of cathepsin L-like proteases in *Trypanosoma rangeli* isolates: markers for diagnosis, genotyping and phylogenetic relationships. *Acta Trop.* 2009;112:249–59.
50. Denise H, Poot J, Jiménez M, Ambit A, Herrmann DC, Vermeulen AN, Coombs GH, et al. Studies on the CPA cysteine peptidase in the *Leishmania infantum* genome strain JPCM5. *BMC Mol Biol.* 2006;7:42.
51. Silva-Almeida M, Pereira BAS, Ribeiro-Guimarães ML, Alves CR. Proteinases as virulence factors in *Leishmania* spp. infection in mammals. *Parasit Vectors.* 2012;5:160.
52. Zampieri RA, Laranjeira-Silva MF, Muxel SM, Lima ACS, Shaw JJ, Floeter-Winter LM. High resolution melting analysis targeting hsp70 as a fast and efficient method for the discrimination of *Leishmania* species. *PLoS Negl Trop Dis.* 2016;10:2.
53. Silva RP, Silva LAMT, Silva MAL, Albuquerque SCG, Goes TCG, Morais RCS, Melo FL, Cavalcante MP. Evaluation of urine for *Leishmania infantum* DNA detection by real-time quantitative PCR. *J Microbiol Methods.* 2016;131:34–41.
54. Costa AP, Ferreira JIGS, Fournier GFSR, Lopes MG, Ramirez D, Acosta ICL, et al. Survey of *Leishmania Infantum Chagasi* in wild and domestic animals in urban area and Atlantic rainforest fragment in northeast, Brazil. *J Biodivers Biopros Dev.* 2014;1:2.
55. Fraga DB, Pacheco LV, Borja LS, Tuy PG, Bastos LA, Solcà MS, Amorim LD, et al. The rapid test based on *Leishmania infantum* chimeric rk28 protein improves the diagnosis of canine visceral Leishmaniasis by reducing the detection of false-positive dogs. *PLoS Negl Trop Dis.* 2016;10:1.
56. Foulet F, Botterel F, Buffet P, Morizot G, Rivollet D, Deniau M, et al. Detection and identification of *Leishmania* species from clinical specimens by using a real-time PCR assay and sequencing of the cytochrome b gene. *J Clin Microbiol.* 2007;45:2110–5.
57. Ceccarelli M, Galluzzi L, Migliazzo A, Magnani M. Detection and characterization of *Leishmania (Leishmania)* and *Leishmania (Viannia)* by SYBR green-based real-time PCR and high resolution melt analysis targeting kinetoplast minicircle DNA. *PLoS One.* 2014;9:2.
58. Ceccarelli M, Galluzzi L, Diotallevi A, Andreoni F, Fowler H, Petersen C, et al. The use of kDNA minicircle subclass relative abundance to differentiate between *Leishmania (L.) infantum* and *Leishmania (L.) amazonensis*. *Parasit Vectors.* 2017;10:1.
59. Dantas-Torres F, Sales KGS, Silva LG, Otranto D, Figueiredo LA. *Leishmania*-FAST15: a rapid, sensitive and low-cost real-time PCR assay for the detection of *Leishmania infantum* and *Leishmania braziliensis* kinetoplast DNA in canine blood samples. *Mol Cell Probes.* 2017;31:65–9.
60. Marfurt J, Niederwieser I, Makia ND, Beck HP, Felger I. Diagnostic genotyping of old and New World *Leishmania* species by PCR-RFLP. *Diagn Microbiol Infect Dis.* 2003;46:115–24.
61. Garcia FB, Santos SSR, Chocay MF, Medeiros ACR, Roselino AMF. Métodos subsidiários para o diagnóstico da Leishmaniose tegumentar americana (LTA): comparação dos resultados do sequenciamento de DNA e da PCR-RFLP para determinação da espécie de *Leishmania* em amostras cutâneo-mucosas. *An Bras.* 2005;80:339–44.
62. Mohammadi MA, Bamorovat M, Harandi MF, Karimi T, Sharifi I, Aflatoonian MR. Comparison of three PCR-based methods for simplicity and cost effectiveness identification of cutaneous Leishmaniasis due to *Leishmania tropica*. *Iran J Parasitol.* 2017;12:215–23.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.