

CORRECTION

Open Access



Correction to: The prevalence of cryptococcal antigen (CrAg) and benefits of pre-emptive antifungal treatment among HIV-infected persons with CD4+ T-cell counts < 200 cells/ μ L: evidence based on a meta-analysis

Yao Li^{1,2†}, Xiaojie Huang^{2†}, Hui Chen³, Yuanyuan Qin^{1,2}, Jianhua Hou², Aixin Li², Hao Wu², Xiaofeng Yan⁴ and Yaokai Chen^{1*}

Correction to: BMC Infect Dis 20, 410 (2020)
<https://doi.org/10.1186/s12879-020-05126-z>

Following publication of the original article [1], the authors identified an error in Fig. 3. The correct figure is given below.

The original article [1] has been corrected.

The original article can be found online at <https://doi.org/10.1186/s12879-020-05126-z>.

* Correspondence: yaokaichen@hotmail.com

†Yao Li and Xiaojie Huang contributed equally to this work.

¹Division of infectious Diseases, Chongqing Public Health Medical Center, 109 Baoyu Road, Shapingba District, Chongqing 400036, China
Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.




Fig. 3 Forest plots of incidence of CM among CrAg + persons receiving azole vs. no intervention or placebo. Abbreviations: M-H, Mantel Haenszel; CI, confidence interval. (“Azole inter” means “Azole drug intervention”, “No inter or plac” mean “No intervention or placebo”

Author details

¹Division of infectious Diseases, Chongqing Public Health Medical Center, 109 Baoyu Road, Shapingba District, Chongqing 400036, China. ²Center for Infectious Diseases, Beijing Youan Hospital, Capital Medical University, Beijing, China. ³School of Biomedical Engineering, Capital Medical University, Beijing, China. ⁴Section of Medical Affairs Administration, Chongqing Public Health Medical Center, Chongqing, China.

Published online: 24 June 2021

Reference

- Li, et al. The prevalence of cryptococcal antigen (CrAg) and benefits of pre-emptive antifungal treatment among HIV-infected persons with CD4+ T-cell counts < 200 cells/μL: evidence based on a meta-analysis. BMC Infect Dis. 2020;20:410. <https://doi.org/10.1186/s12879-020-05126-z>.